

# Chapter 3: Second Order ODE

## §3.1 Introduction to Second Order ODE

Satya Mandal, KU

10 February 2018

# Second Order DE

- ▶ For many, the first encounter with second order ODE occurs, as one starts getting familiar with the concept of acceleration. Recall, acceleration is  $\frac{d^2y}{dt^2}$  where  $y$  is distance travelled. Second Order ODE (SODE) has wide range of applications, in the undergraduate course in Physics and Engineering, for the same reason.
- ▶ SODE models are used in fluid dynamics, heat equations, wave motion, economics and so on.
- ▶ More importantly, a wide variety of SODEs can be solved by analytically.

# Definition: SODEs

**Definition.** A second order ODE has the form

$$\frac{d^2y}{dt^2} = f(t, y, y') \quad (1)$$

where  $f$  is a function of  $t, y, y' := \frac{dy}{dt}$ .

# Linear SODEs

- ▶ A SODE (1) is called a linear SODE (**LSODE**), if  $f$  is linear. That means, if DE (1) has the form:

$$\frac{d^2y}{dt^2} = f\left(t, y, \frac{dy}{dt}\right) = g(t) - q(t)y - p(t)\frac{dy}{dt}$$

where  $g(t)$ ,  $p(t)$ ,  $q(t)$  are functions of  $t$ .

# Standard form of LSOEs

Such LSOEs are often written in the following forms:

- ▶ First,

$$\frac{d^2y}{dt^2} + p(t)\frac{dy}{dt} + q(t)y = g(t) \quad (2)$$

where  $p(t)$ ,  $q(t)$ ,  $g(t)$  are functions of  $t$ . Then, also as:

$$P(t)\frac{d^2y}{dt^2} + Q(t)\frac{dy}{dt} + R(t)y = G(t) \quad (3)$$

where  $P(t)$ ,  $Q(t)$ ,  $R(t)$ ,  $G(t)$  are functions of  $t$ .

- ▶ The LSOE (3) can be reduced to (2), when  $P(t) \neq 0$ , and conversely.

# Recall and Compare:

Recall the form of the first order linear ODE and compare:

$$\begin{cases} 1^{\text{st}} \text{ Order Linear : } \frac{dy}{dt} + p(t)y = g(t) \\ 2^{\text{nd}} \text{ Order Linear : } \frac{d^2y}{dt^2} + p(t)\frac{dy}{dt} + q(t)y = g(t) \end{cases}$$

# Initial value problems in SODEs

- ▶ An initial value problem (IVP) in SODE consists of ODE (1), (2), or (3) together with initial value conditions:

$$y(t_0) = y_0, \quad y'(t_0) = y'_0.$$

So, one such form of a second order IVP is:

$$\begin{cases} \frac{d^2y}{dt^2} + p(t)\frac{dy}{dt} + q(t)y = g(t) \\ y(t_0) = y_0, \quad y'(t_0) = y'_0. \end{cases} \quad (4)$$

- ▶ As a rule of Thumb, two equations are needed to determine two unknowns. It would be evident later, that general solutions of (1) involve two arbitrary constants  $c_1, c_2$ . That is (4) has two conditions.

# Homogeneous LSODEs

- ▶ A LSODE is called **homogeneous**, if  $g(t) = 0$  in (2), or if  $G(t) = 0$  in (3).
- ▶ If  $g(t) \neq 0$  [resp.  $G(t) \neq 0$ ], the equations (2) [resp. (3)] would be called a **nonhomogeneous** linear equation.
- ▶ So, the a homogeneous LSODE can be written as

$$P(t)\frac{d^2y}{dt^2} + Q(t)\frac{dy}{dt} + R(t)y = 0 \quad (5)$$

- ▶ In fact (§ 3.5?), solutions of homogeneous LSODE (5) leads to solutions of nonhomogeneous LSODE (3).



# Rest of this Chapter

- ▶ In next few sections, we solve homogeneous equations (5) with constant coefficients. So, they look like:

$$a \frac{d^2 y}{dt^2} + b \frac{dy}{dt} + cy = 0 \quad a, b, c \in \mathbb{R}. \quad (6)$$

However, we comment on ODEs (5), as is.

- ▶ In latter sections, we solve nonhomogeneous equations, whose left side is as in (6). So, they look like:

$$a \frac{d^2 y}{dt^2} + b \frac{dy}{dt} + cy = g(t) \quad (7)$$