

On Dévissage for Witt groups

Satya Mandal¹ and Sarang Sane²

¹*University of Kansas, Lawrence, Kansas 66045*
e-mail: mandal@ku.edu

²*IIT Madras, Sardar Patel Road, Adyar, Chennai 600 036*
e-mail: sarangsanemath@gmail.com

Communicated by: Bangere P. Purnaprajna

Received: August 17, 2015

Abstract. In this paper we extend and apply the work of Paul Balmer and others on derived and triangular Witt Groups. We obtain a generalized form of dévissage for derived Witt Groups over Cohen-Macaulay rings.

2000 *Mathematics Subject Classification:* Primary: 11E81, 18E30, 19G12; Secondary: 13C10, 13D

1. Introduction

This is the first one of a number of articles ([MS,M1,M2]) dedicated to extending the theory of derived Witt groups, due to Paul Balmer, to non-regular schemes. We restrict ourselves to the derived categories of complexes, whose homologies have finite projective dimension. This introduction also serves as a prelude to the same for all the articles in this series.

In this article, we develop some background for this series, including a definition of Witt groups of full subcategories of triangulated categories with duality, and a version of the dévissage theorem [BW, Theorem 6.1] for Witt groups of Cohen-Macaulay rings. To introduce this theorem, let A be a commutative Noetherian domain of dimension d with 2 invertible and K be its quotient field. It is a classical question (known as purity) as to when the map $W(A) \rightarrow W(K)$ is injective. Purity is a conjecture when A is a regular local ring and is affirmatively settled in ([CS,OP,OPSS]).

In general, we can extend the above map to the right for any regular scheme by considering the Gersten-Witt complex. Let $X = \text{Spec}(A)$ be of

Partially supported by a General Research Grant from KU.

dimension d with 2 invertible and $X^{(n)}$ denote the points of codimension n . A Gersten-Witt complex

$$0 \rightarrow W(A) \rightarrow \bigoplus_{x \in X^{(0)}} W(k(x)) \rightarrow \bigoplus_{x \in X^{(1)}} W(k(x)) \rightarrow \dots \rightarrow \bigoplus_{x \in X^{(d)}} W(k(x)) \rightarrow 0$$

was first constructed by Pardon [Pa1]. He further conjectured the exactness of this sequence for regular local rings and affirmatively settled it in many cases.

Subsequently, with the introduction of triangular Witt groups by Balmer [B1,B2,B3], Witt groups could be viewed as a cohomology theory. Using this, another Gersten-Witt complex could be defined (though both complexes look similar, it seems unproven that the differentials match) [BW], similar to the one in K-theory. Once again it is an open question as to when the complex is exact. In particular, it is conjectured to be so when $X = \text{Spec}(A)$ where A is a regular, local ring and this is affirmatively settled in ([B4,Pa1,Pa2,B4,BGPW]). [B1,B2,B3,BW] form the basic backbone of the methodologies in this article and we would like to remark that the interested reader would be well advised to take a look at them. *For any unexplained notations and definitions in the introduction, please refer to (2.1).*

The key result which allows one to move from derived Witt groups to the Gersten-Witt complex is dévissage [BW, Section 6] which states that for a regular, local ring (A, \mathfrak{m}, k) , of dimension d , we have $W^n(D_{fl}^b(A)) \cong W(k)$ if $n \equiv d \pmod 4$ and $W^n(D_{fl}^b(A)) \cong 0$ otherwise. We describe below the generalized form of this theorem for a Cohen-Macaulay ring A with $\dim A_{\mathfrak{m}} = d$ for all maximal ideals \mathfrak{m} and 2 invertible.

Suppose A is a Cohen-Macaulay ring with $\dim A = d$. Since there are modules of infinite projective dimension over such a ring and of finite length, the usual duality $Ext^d(_, A)$ does not work well. The options are either to change the duality (w.r.t. the canonical module) but then use the category of all modules (coherent Witt groups) or impose finite projective dimension homology conditions on the complexes. The first path is taken and deeply studied in Pardon [Pa2,Pa3] and more recently in Gille [G1,G2], where they also establish a Gersten-Witt complex of coherent Witt groups.

In this article, we take the second approach. Let $\mathcal{MFPD}(A)$ be the category of finitely generated A -modules with finite projective dimension, $\mathcal{MFL}(A)$ be the category of finitely generated A -modules with finite length, and $\mathcal{A} = \mathcal{MFPD}_{fl}(A)$ be the full subcategory of finitely generated A -modules with finite projective dimension and finite length (the “intersection”). Note then that \mathcal{A} is an exact category and has a natural duality given by $M \mapsto Ext_A^d(M, A)$ and so we can consider the Witt group $W(\mathcal{A})$. By Balmer [B2], we already know that $W(\mathcal{A}) \cong W^d(D^b(\mathcal{A}))$.

We consider the category $D_{\mathcal{A}}^b(\mathcal{A})$ with homologies in \mathcal{A} . Based on the fact that \mathcal{A} actually has the 2-out-of-3 property for objects, we prove that the

duality actually restricts to $D_{\mathcal{A}}^b(\mathcal{A})$ and this allows a suitable definition of Witt groups $W^i(D_{\mathcal{A}}^b(\mathcal{A}))$. Once defined, we prove that the above isomorphism actually factors through isomorphisms $W(\mathcal{A}) \xrightarrow{\sim} W^d(D_{\mathcal{A}}^b(\mathcal{A})) \xrightarrow{\sim} W^d(D^b(\mathcal{A}))$.

We now consider the category $D_{\mathcal{A}}^b(\mathcal{P}(A))$. Similar to $D_{\mathcal{A}}^b(\mathcal{A})$, we establish that this category is stable under duality and define the Witt groups $W^i(D_{\mathcal{A}}^b(\mathcal{A}))$. Having done so, we prove our version of dévissage (5.12,6.7):

$$\begin{aligned} W(\mathcal{A}) &\xrightarrow{\sim} W^d(D_{\mathcal{A}}^b(\mathcal{P}(A))) \\ W^-(\mathcal{A}) &\xrightarrow{\sim} W^{d+2}(D_{\mathcal{A}}^b(\mathcal{P}(A))) \\ W^{d+1}(D_{\mathcal{A}}^b(\mathcal{P}(A))) &\cong W^{d-1}(D_{\mathcal{A}}^b(\mathcal{P}(A))) \cong 0. \end{aligned}$$

Note that when A is regular, this is exactly the same theorem as that in [BW]. Further, in the Gorenstein case, there is a natural commuting square involving the above dévissage statement and the dévissage statement in [G1], for coherent Witt groups. As of now, we do not know if these sets of groups coincide or not, which would also be a subject of future investigations. However, we point out that there are more forms and neutral spaces in the realm of coherent Witt theory. Since we prove the theorem without regularity assumptions or the existence of a natural dualizing complex, we do not have access to the equivalences of the derived categories with duality as in [BW] or [G1] (in particular we cannot use the powerful lemma of Keller [K, §1.5, Lemma and Example(b)]). Our proof is thus necessarily more elementary and naïve than the one in [BW]. One of the key ingredients in the proof is the construction of a special sublagrangian (5.6) for symmetric forms in $(D_{\mathcal{A}}^b(\mathcal{P}(A)))$.

In deed, the methods in this paper have much wider applications and set the tone of arguments used in the rest of the articles in this series (*loc. cit.*). A key to these methods is to use the sublagrangian theorem of Balmer ([B3]), to construct symmetric forms of complexes, Witt equivalent to a given form, with shorter length. Methods also provide further insight into nonsingular varieties ([M1]). Our interest in these studies stems from the introduction of the Chow-Witt groups $\widehat{CH}^r(A)$ for $0 \leq r \leq d$, due to Barge and Morel [BM] and developed by Fasel [F1], as the obstruction groups for splitting of projective modules, which works best for nonsingular varieties. This also serves as a motivation for our interest in maintaining the category of projective modules in our statement of dévissage. Jean Fasel informs that, for singular varieties X , Chow-Witt groups $\widehat{CH}^r(X)$ and obstruction classes can be defined in the same manner, using coherent Witt groups. However, it is not known whether vanishing of the obstruction classes would lead to splitting. We feel that, for the purpose of developing an obstruction theory for singular varieties, it would be more natural to consider some analogue of the derived

Witt groups of the category of projective modules. This approach would be of its own independent interest for both computations and applications.

We make a few comments about the layout of the article. The article was written with the readership in mind and provides some extra details. In section 2, we establish the basic definitions and a key result on projective dimensions. In the section 3, we establish the important theorem that the categories $D_{\mathcal{A}}^b(\mathcal{P}(A))$ and $D_{\mathcal{A}}^b(\mathcal{A})$ are closed under duality, and more specifically how the homologies of the dual look like. Once this is established, in section 4, we define the Witt groups of the above categories and expectedly, they are 4-periodic, i.e. $W^n(D_{\mathcal{A}}^b(\mathcal{P}(A))) \xrightarrow{\sim} W^{n+4}(D_{\mathcal{A}}^b(\mathcal{P}(A)))$. Finally, in sections 5 and 6, we prove our main theorems about dévissage.

Acknowledgements

We would like to thank Sankar P. Dutta for many helpful discussions, and particularly, for example (2.4). We are also very thankful to Jean Fasel for many helpful discussions and guidance. Thanks are also due to Manuel Ojanguren for some helpful corrections and comments. The second named author is thankful to the University of Kansas and the Robert D. Adams Trust.

2. Basic notations and preliminaries

Notations 2.1. Throughout this article, A will denote a Cohen-Macaulay ring with $\dim A_m = d \geq 2$, for all maximal ideals m of A . Further, 2 is always invertible in A . We set up the notations:

1. $\mathcal{M}(A)$: category of finitely generated A -modules.
2. $\mathcal{MFPD}(A)$: full subcategory of finitely generated A -modules with finite projective dimension.
3. $\mathcal{MFL}(A)$: category of finitely generated A -modules with finite length.
4. $\mathcal{A} = \mathcal{MFPD}_{fl}(A)$: category of finitely generated A -modules with finite length and finite projective dimension.
5. $\mathcal{P}(A)$: category of finitely generated projective A -modules.
6. For any exact category \mathcal{C} , $Ch^b(\mathcal{C})$ is the category of bounded chain complexes with objects in \mathcal{C} , and $D^b(\mathcal{C})$ is its derived category.
7. For any two exact categories \mathcal{C}, \mathcal{D} in an ambient abelian category \mathcal{C}' , $Ch_{\mathcal{D}}^b(\mathcal{C})$ is the full subcategory of $Ch^b(\mathcal{C})$ consisting of complexes with homologies in \mathcal{D} . $D_{\mathcal{D}}^b(\mathcal{C})$ is its derived category, which is also the full subcategory of $D^b(\mathcal{C})$ consisting of objects from $Ch_{\mathcal{D}}^b(\mathcal{C})$.
8. \mathcal{R} : full subcategory of $D_{\mathcal{A}}^b(\mathcal{P}(A))$ consisting of objects P_{\bullet} such that $P_i = 0$ for $i > d, i < 0$ and $H_i(P_{\bullet}) = 0$ for all $i \neq 0$ and $H_0(P_{\bullet}) \in \mathcal{A}$.
9. For objects M in \mathcal{A} , let $M^{\vee} = Ext_A^d(M, A)$ and $\tilde{w} : M \xrightarrow{\sim} M^{\vee\vee}$ be the identification by double ext. (but we make this more precise in diagram ((5)) and the explanation of i .)

10. We will denote complexes P_\bullet by:

$$\cdots 0 \longrightarrow P_m \xrightarrow{\hat{c}_m} P_{m-1} \longrightarrow \cdots \longrightarrow P_n \longrightarrow 0 \cdots$$

- 11. A non-zero complex P_\bullet is said to be supported on $[m, n]$ if $P_i = 0$ for all $i < m$ and $i > n$.
- 12. For a complex P_\bullet of projective A -modules P_\bullet^* will denote the usual dual induced by $\text{Hom}(*, A)$ and $\varpi : P_\bullet \xrightarrow{\sim} P_\bullet^{**}$ will denote the identification by evaluation. Note that the degree r -component of the dual P_\bullet^* is $(P_{-r})^*$.
- 13. The **length** of a non-zero complex P_\bullet is defined as $\ell(P_\bullet) = u - l$ where $P_u \neq 0, P_l \neq 0$ and $P_i = 0$ for all $i < l$ and $i > u$.
- 14. Let $B_r = B_r(P_\bullet) := \hat{\partial}_{r+1}(P_{r+1}) \subseteq P_r$ denote the module of r -boundaries and $Z_r = Z_r(P_\bullet) := \ker(\hat{\partial}_r) \subseteq P_r$ denote the module of r -cycles (or the r^{th} syzygy).
- 15. The r^{th} -homology of P_\bullet will be denoted by $H_r = H_r(P_\bullet) := \frac{Z_r}{B_r}$.
So, the r^{th} -homology of the dual is $H_r(P_\bullet^*) = \frac{\ker(\hat{\partial}_{-(r-1)}^*)}{\text{image}(\hat{\partial}_r^*)}$.
- 16. A full exact subcategory \mathcal{C} of an abelian category \mathcal{D} is said to have the 2-out-of-3 property if for every short exact sequence in \mathcal{D} , whenever two of the objects are objects of \mathcal{C} , then so is the third.

Remark 2.2. The subcategory $\mathcal{MFPD}(A)$ and \mathcal{A} are both exact subcategories and in fact have the 2-out-of-3 property. The category \mathcal{R} is also an exact category. Although it is a subcategory of $D_{\mathcal{A}}^b(\mathcal{P}(A))$, it has no translation and is actually naturally equivalent to the category \mathcal{A} . The natural functor $\eta : \mathcal{R} \rightarrow \mathcal{A}$ is given by sending a complex Q_\bullet to $H_0(Q_\bullet)$. The inverse functor ι is given by associating to objects $M \in \mathcal{A}$ a projective resolution of length d .

Note further that when A is not regular, the categories $D_{\mathcal{A}}^b(\mathcal{P}(A))$ and $D_{\mathcal{MFPD}(A)}^b(\mathcal{P}(A))$ are not closed under the cone operation as the following example demonstrates.

Example 2.3. Let (A, \mathfrak{m}) be a non-regular Cohen-Macaulay ring with $\dim A = d$, such that $\mathfrak{m} = (f_1, f_2, \dots, f_d, z)$. We can assume, using prime avoidance, that f_1, f_2, \dots, f_d is a regular sequence. Let $U_\bullet = \text{Kos}_\bullet(f_1, f_2, \dots, f_d)$ be the Koszul complex. Since the only nonzero homology of U_\bullet is $H_0(U_\bullet) = \frac{A}{(f_1, f_2, \dots, f_d)} \in \mathcal{A}$, U_\bullet and all its translates are objects of both the categories above. Let $C(z)$ denote the cone of the chain complex map $z : U_\bullet \rightarrow U_\bullet$. From the long exact homology sequence corresponding to the short exact sequence of chain complexes

$$0 \longrightarrow U_\bullet \longrightarrow C(z) \longrightarrow U_\bullet[1] \longrightarrow 0$$

it follows that

$$H_0(\text{Cone}(z)) \cong \text{coker} \left(\frac{A}{(f_1, f_2, \dots, f_d)} \xrightarrow{z} \frac{A}{(f_1, f_2, \dots, f_d)} \right) \cong \frac{A}{\mathfrak{m}} \notin \mathcal{A}.$$

So, $C(z)$ is not an object in $D_{\mathcal{A}}^b(\mathcal{P}(A))$.

Next, we ask if $Ch_{\mathcal{MFPD}}^b(\mathcal{P}(A))$ is closed under duality. We thank Sankar Dutta for providing the following example:

Example 2.4 (Dutta). Let (A, \mathfrak{m}, k) be any non-regular Cohen-Macaulay local ring, with $\dim A = d$. Let

$$\cdots \longrightarrow P_d \xrightarrow{\partial_d} P_{d-1} \longrightarrow \cdots \longrightarrow P_0 \longrightarrow k \longrightarrow 0$$

be a projective resolution of k . Let $*$ denote $Hom(-, A)$ and $M = \text{cokernel}(\partial_d^*)$. Since $Ext^r(k, A) = 0$ for all $0 \leq r < d$, the sequence

$$0 \longrightarrow P_0^* \longrightarrow \cdots \longrightarrow P_{d-1}^* \longrightarrow P_d^* \longrightarrow M \longrightarrow 0$$

is a projective resolution of M . Dualizing this sequence, it follows that $Ext_A^d(M, A) \cong k$, which does not have finite projective dimension. In particular, $Ch_{\mathcal{MFPD}}^b(\mathcal{P}(A))$ is not closed under duality.

Note however that in the above example, M does not have finite length. Indeed, we will prove in section 3 that the category $Ch_{\mathcal{A}}^b(\mathcal{P}(A))$ is closed under duality.

We mention a few standard results for the sake of completeness.

Lemma 2.5. *Let (A, \mathfrak{m}) be a Cohen-Macaulay local ring with $\dim A = d$. Let $M \in \mathcal{MFL}(A)$. Then Further, $Ext^i(M, A) = 0$ for all $i < d$. and $Ext^d(M, A) \neq 0$ is also in $\mathcal{MFL}(A)$. Note further that if $M \in \mathcal{A}$, then so is $Ext^d(M, A)$.*

Lemma 2.6. *Let A be a Cohen-Macaulay ring with $\dim A = d$. Let $M \in \mathcal{A}$. There is a natural isomorphism*

$$\varpi : M \xrightarrow{\sim} M^{\vee\vee}.$$

Corollary 2.7. *$(\mathcal{A}^{\vee}, \pm \tilde{\varpi})$ are exact categories with duality.*

Proposition 2.8. *Let A be Cohen-Macaulay with $d = \dim A_{\mathfrak{m}} \geq 2$ for all maximal ideals \mathfrak{m} . Let P_{\bullet} be a complex in $Ch^b(\mathcal{P}(A))$. Assume that all the homologies $H_r := H_r(P_{\bullet}) \in \mathcal{A}$. Then we claim:*

1. *The modules B_r and Z_r have finite projective dimension $\forall r$. In that case, $\text{proj dim}(Z_r) = \text{proj dim}(B_{r-1}) - 1$.*

2. For all $r \in \mathbb{Z}$, we have $\text{proj dim } B_r \leq d - 1$ and so $\text{proj dim } Z_r \leq d - 2$.
3. If $H_r \neq 0$ then $\text{proj dim } B_r = d - 1$.

Proof. Note that P_\bullet is a bounded complex and so let it be supported on $[m, n]$. Then $Z_n = P_n$ and so has finite projective dimension. Since there are short exact sequences:

$$0 \rightarrow B_r \rightarrow Z_r \rightarrow H_r \rightarrow 0 \quad 0 \rightarrow Z_r \rightarrow P_r \rightarrow B_{r-1} \rightarrow 0,$$

it is clear that if Z_r is of finite projective dimension, then so are B_r and Z_{r+1} , and hence the proof follows by induction. The second part of (1) also follows from the above exact sequence.

Since B_r is torsion free, it has depth at least 1 and hence, by the Auslander-Buchsbaum theorem, $\text{proj dim}(B_r) \leq d - 1$. So, $\text{proj dim}(Z_r) \leq d - 2$. So, (2) is established.

Now, assume $H_r \neq 0$. Choose a maximal ideal \mathfrak{m} in the support of H_r . Then, consider the localized short exact sequence

$$0 \rightarrow (B_r)_{\mathfrak{m}} \rightarrow (Z_r)_{\mathfrak{m}} \rightarrow (H_r)_{\mathfrak{m}} \rightarrow 0.$$

Then, we get a long exact sequence of $\text{Tor}_{A_{\mathfrak{m}}}(-, A/\mathfrak{m})$, which gives us that

$$\text{Tor}_{A_{\mathfrak{m}}}^d((H_r)_{\mathfrak{m}}, A/\mathfrak{m}) \cong \text{Tor}_{A_{\mathfrak{m}}}^{d-1}((B_r)_{\mathfrak{m}}, A/\mathfrak{m}) \quad \text{and} \quad \text{Tor}_{A_{\mathfrak{m}}}^d((B_r)_{\mathfrak{m}}, A/\mathfrak{m}) \cong 0,$$

since we have already proved that $\text{proj dim}_A Z_r \leq d - 2$. Thus we obtain $\text{proj dim}_{A_{\mathfrak{m}}}(B_r)_{\mathfrak{m}} = d - 1$. Since we know that $\text{proj dim}_A(B_r) \leq d - 1$, this implies $\text{proj dim}_A(B_r) = d - 1$. This establishes (3). \square

The complexes in $\text{Ch}^b(\mathcal{P}(A))$ with finite length homologies have at least d nonzero components at the left where the homology is 0. This proposition plays a key role in sections 5 and 6.

Proposition 2.9. *Let A be Cohen-Macaulay with $d = \dim A_{\mathfrak{m}}$ for all maximal ideals \mathfrak{m} . Let P_\bullet be a bounded complex of projective modules, such that $H_i = 0 \forall i > n$ and $H_n \neq 0$ is of finite length. Then $P_i \neq 0, n \leq i \leq n + d$.*

Proof. Since $H_i = 0 \forall i > n$ and the complex is bounded, we get that $\frac{P_n}{B_n}$ is of finite projective dimension, since the components with indices $\geq n$ give a resolution. Now, let \mathfrak{m} be a maximal ideal in the support of $H_n(P_\bullet)$, then $(H_n(P_\bullet))_{\mathfrak{m}} \subseteq \frac{(P_n)_{\mathfrak{m}}}{(B_n)_{\mathfrak{m}}}$ is of finite length, and hence $\frac{(P_n)_{\mathfrak{m}}}{(B_n)_{\mathfrak{m}}}$ has depth 0. By the Auslander-Buchsbaum theorem, $\text{proj dim}_{A_{\mathfrak{m}}} \left(\frac{(P_n)_{\mathfrak{m}}}{(B_n)_{\mathfrak{m}}} \right) = d$. But that means $\text{proj dim}_A \left(\frac{P_n}{B_n} \right) = d$. Hence, the resolution of $\frac{P_n}{B_n}$ given by the components of P_\bullet with indices $\geq n$ must have length at least d . Hence, $P_i \neq 0, n \leq i \leq n + d$. The proof is complete. \square

3. Duality

As always, A will denote a Cohen-Macaulay ring with $\dim A_m = d \geq 2$, for all maximal ideals m and $\mathcal{A} = \mathcal{MFPD}_{fl}(A)$. In this section, we prove that the category $Ch_{\mathcal{A}}^b(\mathcal{P}(A))$ is closed under duality and give a precise description of the homologies of the dual.

Theorem 3.1. *Suppose P_{\bullet} is a complex in $Ch^b(\mathcal{P}(A))$ with homologies in \mathcal{A} . Then we have:*

$$Ext^i(Z_r, A) \cong \begin{cases} Ext^d(H_{r+i-(d-1)}, A) & 1 \leq i \leq d-2 \\ 0 & \text{for } i \geq d-1 \end{cases} \quad (1)$$

$$Ext^i(B_r, A) \cong \begin{cases} Ext^d(H_{r+i-(d-1)}, A) & 1 \leq i \leq d-1 \\ 0 & \text{for } i \geq d \end{cases} \quad (2)$$

$$Ext^i\left(\frac{P_r}{B_r}, A\right) \cong \begin{cases} Ext^d(H_{r+i-d}, A) & 1 \leq i \leq d \\ 0 & i \geq d \end{cases} \quad (3)$$

Proof. Since $P_i = 0 \forall i \ll 0$, the theorem is true for $r \ll 0$. So, we assume that the theorem is true for $r-1$ and prove it for r .

Corresponding to the short exact sequence $0 \rightarrow Z_r \rightarrow P_r \rightarrow B_{r-1} \rightarrow 0$, we get a long exact Ext-sequence which yields

$$0 \rightarrow Ext^0(B_{r-1}, A) \rightarrow P_r^* \rightarrow Ext^0(Z_r, A) \rightarrow Ext^1(B_{r-1}, A) \rightarrow 0 \quad (4)$$

and for $i \geq 1$ we have $Ext^i(Z_r, A) \cong Ext^{i+1}(B_{r-1}, A)$. Thus, the induction hypothesis yields that for $i \geq 1$,

$$Ext^i(Z_r, A) = Ext^{i+1}(B_{r-1}, A) = \begin{cases} Ext^d(H_{r+i-(d-1)}, A) & 1 \leq i \leq d-2 \\ 0 & \text{for } i \geq d-1 \end{cases}$$

So, equation (1) is established.

Consider the long exact Ext-sequence corresponding to the short exact sequence $0 \rightarrow B_r \rightarrow Z_r \rightarrow H_r \rightarrow 0$. By (2.5), $Ext^i(H_r, A) = 0$ for all $i \neq d$ and since $Ext^i(Z_r, A) = 0 \forall i > d-2$ from equation (1), it follows that

$$\begin{aligned} Ext^i(B_r, A) &\cong \begin{cases} Ext^i(Z_r, A), & 0 \leq i \leq d-2 \\ Ext^d(H_r, A), & i = d-1 \\ 0, & i \geq d \end{cases} \\ &\cong \begin{cases} Ext^0(Z_r, A), & i = 0 \\ Ext^d(H_{r+i-(d-1)}, A), & 1 \leq i \leq d-1 \\ 0, & i \geq d \end{cases} \end{aligned}$$

Now consider the short exact sequence $0 \rightarrow H_r \rightarrow \frac{P_r}{B_r} \rightarrow B_{r-1} \rightarrow 0$. Again, $\text{Ext}^i(H_r, A) = 0 \forall i \neq d$ from (2.5) and from equation (2) we get $\text{Ext}^i(B_r, A) = 0 \forall i > d - 1$. So it follows that

$$\begin{aligned} \text{Ext}^i\left(\frac{P_r}{B_r}, A\right) &\cong \begin{cases} \text{Ext}^i(B_{r-1}, A), & 0 \leq i \leq d - 1 \\ \text{Ext}^d(H_r, A), & i = d \\ 0, & i > d \end{cases} \\ &\cong \begin{cases} \text{Ext}^i(B_{r-1}, A), & i = 0 \\ \text{Ext}^d(H_{r+i-d}, A), & 1 \leq i \leq d \\ 0, & i > d \end{cases} \end{aligned}$$

□

Corollary 3.2. *Suppose P_\bullet is a complex in $\text{Ch}^b(\mathcal{P}(A))$ with homologies in \mathcal{A} . Then, for all $r \in \mathbb{Z}$*

$$\text{Ext}^i(B_r, A), \quad \text{Ext}^i(Z_r, A), \quad \text{Ext}^i\left(\frac{P_r}{B_r}, A\right)$$

are in \mathcal{A} for $i \geq 1$ and are in $\mathcal{MFPD}(A)$ for $i = 0$.

Proof. By (2.5) and the preceding theorem (3.1), for $i \geq 1$, the statement is clear. For $i = 0$, we recall below equation (4) from the preceding proof:

$$0 \rightarrow \text{Ext}^0(B_{r-1}, A) \rightarrow P_r^* \rightarrow \text{Ext}^0(Z_r, A) \rightarrow \text{Ext}^1(B_{r-1}, A) \rightarrow 0.$$

and that we also proved $\text{Ext}^0(B_r, A) \cong \text{Ext}^0(Z_r, A)$ and $\text{Ext}^0\left(\frac{P_r}{B_r}, A\right) \cong \text{Ext}^0(B_{r-1}, A)$. Hence, it is enough to know that $\text{Ext}^0(B_{r-1}, A)$ satisfies the theorem. Once again induction saves the day! □

This allows us to conclude our main theorem of the section.

Theorem 3.3. *Let P_\bullet be a complex as in theorem (3.1). Then, for $t \in \mathbb{Z}$, we have*

$$H_{-t}(P_\bullet^*) \cong \text{Ext}^d(H_{t-d}(P_\bullet), A) \cong H_{t-d}(P_\bullet)^\vee.$$

In particular, $H_r(P_\bullet^) \in \mathcal{A}$ and hence, $\text{Ch}_\mathcal{A}^b(\mathcal{P}(A))$ is closed under duality.*

Proof. Consider the dual complex: $\dots P_{t-1}^* \xrightarrow{(\partial_t)^*} P_t^* \xrightarrow{(\partial_{t+1})^*} P_{t+1}^* \dots$. Note that $(\partial_{t+1})^* : P_t^* \rightarrow P_{t+1}^*$ factors through

$$P_t^* \rightarrow B_t^* \hookrightarrow \left(\frac{P_{t+1}}{B_{t+1}}\right)^* \hookrightarrow P_{t+1}^*$$

(recall $d \geq 2$) and hence, $\ker((\partial_{t+1})^*) = \ker(P_t^* \rightarrow B_t^*) = \left(\frac{P_t}{B_t}\right)^*$. Similarly, $\ker((\partial_t)^*) = \left(\frac{P_{t-1}}{B_{t-1}}\right)^*$ and hence we obtain the exact sequence

$$0 \rightarrow \left(\frac{P_{t-1}}{B_{t-1}}\right)^* \rightarrow (P_{t-1})^* \rightarrow \left(\frac{P_t}{B_t}\right)^* \rightarrow H_{-t}(P_\bullet^*) \rightarrow 0.$$

Note also that there is an exact Ext-sequence

$$0 \rightarrow \left(\frac{P_{t-1}}{B_{t-1}}\right)^* \rightarrow (P_{t-1})^* \rightarrow (B_{t-1})^* \rightarrow \text{Ext}^1\left(\frac{P_{t-1}}{B_{t-1}}, A\right) \rightarrow 0.$$

But since $d \geq 2$, we have

$$\begin{array}{ccccccc} 0 & \longrightarrow & \left(\frac{P_{t-1}}{B_{t-1}}\right)^* & \longrightarrow & (P_{t-1})^* & \longrightarrow & (B_{t-1})^* \longrightarrow \text{Ext}^1\left(\frac{P_{t-1}}{B_{t-1}}, A\right) \longrightarrow 0 \\ & & \parallel & & \parallel & & \downarrow \wr \\ 0 & \longrightarrow & \left(\frac{P_{t-1}}{B_{t-1}}\right)^* & \longrightarrow & (P_{t-1})^* & \longrightarrow & \left(\frac{P_t}{B_t}\right)^* \longrightarrow H_{-t}(P_\bullet^*) \longrightarrow 0 \end{array}$$

Hence, by (3.1), we get that $H_{-t}(P_\bullet^*) \cong \text{Ext}^d(H_{t-d}(P_\bullet), A)$. The rest follows from (2.5). □

Remark 3.4. It is a straightforward diagram check that all the isomorphisms in (3.1) and (3.3) are natural. In particular, that means that if we have a morphism of complexes $P_\bullet \xrightarrow{f} Q_\bullet$, then there is a commutative diagram:

$$\begin{array}{ccc} H_{-t}(Q_\bullet^*) & \xrightarrow{\sim H_{-t}(f^*)} & H_{-t}(P_\bullet^*) \\ \downarrow \wr & & \downarrow \wr \\ H_{t-d}(Q_\bullet)^\vee & \xrightarrow{\sim H_{t-d}(f)^\vee} & H_{t-d}(P_\bullet)^\vee \end{array}$$

Finally, as an easy consequence of the above theorem, we obtain (for free!) that $D_{\mathcal{A}}^b(\mathcal{A})$ is closed under duality $M^\vee = \text{Ext}^d(M, A)$.

Theorem 3.5. *The category $Ch_{\mathcal{A}}^b(\mathcal{A})$ is closed under the duality $^\vee$ induced by the duality $^\vee$ in \mathcal{A} .*

Proof. Suppose M_\bullet is a complex in $Ch_{\mathcal{A}}^b(\mathcal{A})$. Without loss of generality, we assume M_\bullet is supported on $[n, 0]$. Each component M_i has a projective resolution of length d , and putting them together with the induced maps, we get a double complex $L_{\bullet\bullet}$, as in the left figure below:

$$\begin{array}{ccccccc} 0 & \rightarrow & \cdots & \rightarrow & \cdots & \rightarrow & 0 \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ L_{\bullet\bullet} = & 0 \rightarrow & L_{1n} \rightarrow L_{1(n-1)} \rightarrow \cdots \rightarrow L_{11} \rightarrow L_{10} \rightarrow 0 & & L'_{\bullet\bullet} = & 0 \rightarrow & L_{(d-1)0}^* \rightarrow L_{(d-1)1}^* \rightarrow \cdots \rightarrow L_{(d-1)(n-1)}^* \rightarrow L_{(d-1)n}^* \rightarrow 0 \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ & 0 \rightarrow & L_{0n} \rightarrow L_{0(n-1)} \rightarrow \cdots \rightarrow L_{01} \rightarrow L_{00} \rightarrow 0 & & & 0 \rightarrow & L_{d0}^* \rightarrow L_{d1}^* \rightarrow \cdots \rightarrow L_{d(n-1)}^* \rightarrow L_{dn}^* \rightarrow 0 \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ M_\bullet = & 0 \rightarrow & M_n \rightarrow M_{n-1} \rightarrow \cdots \rightarrow M_1 \rightarrow M_0 \rightarrow 0 & & M_\bullet^\vee = & 0 \rightarrow & M_0^\vee \rightarrow M_1^\vee \rightarrow \cdots \rightarrow M_{n-1}^\vee \rightarrow M_n^\vee \rightarrow 0 \end{array}$$

Dualizing, $L'_{\bullet\bullet}$ gives a similar resolution of M_{\bullet}^{\vee} , as shown on the right above (note that there are sign conventions on the differentials of the complexes here, in particular M_{\bullet}^{\vee} acquires a $(-1)^d$ factor on its differentials. We refer to [W] for the conventions for total complexes.)

Now, the total complexes give quasi-isomorphisms

$$Tot(L_{\bullet\bullet}) \longrightarrow M_{\bullet}, \quad Tot(L'_{\bullet\bullet}) \twoheadrightarrow M_{\bullet}^{\vee}.$$

So, $H_i(Tot(L_{\bullet\bullet})) \in \mathcal{A}$ for all $i \in \mathbb{Z}$. By (3.3), $H_i(Tot(L'_{\bullet\bullet})) \in \mathcal{A}$ for all $i \in \mathbb{Z}$. Now after translating $Tot(L_{\bullet\bullet})^*$ d components to the left, we observe that it is actually chain homotopy equivalent to $Tot(L'_{\bullet\bullet})$ and so we have $H_i(Tot(L'_{\bullet\bullet})) \in \mathcal{A}$. Finally, the above quasi-isomorphism yields that $H_i(M_{\bullet}^{\vee}) \xrightarrow{\sim} H_i(Tot(L'_{\bullet\bullet})) \in \mathcal{A}$. This completes the proof. The proof is complete. \square

4. Definitions of Witt groups

In this section, we define Witt groups of the categories we work with. In particular, we extend the definition of Witt groups from triangulated categories with duality to their additive subcategories which are closed under orthogonal sums, translations and isomorphisms. Since it is possible that there is cause for confusion about translation, we start by clearing the air.

Definition 4.1. In all the categories of complexes, there are two possible translations, T_u and T_s . The complex $T_u P_{\bullet}$ is defined as $(T_u P_{\bullet})_i = P_{i-1}$ and $\partial(T_u P_{\bullet})_i = \partial(P_{\bullet})_{i-1}$. The complex $T_s P_{\bullet}$ is defined as $(T_s P_{\bullet})_i = P_{i-1}$ and $\partial(T_s P_{\bullet})_i = -\partial(P_{\bullet})_{i-1}$.

Note that T_s seems to be the “standard” translation in literature and that is always the translation we use on any category of complexes.

However, given a duality $*$ on such a category (e.g. $D_{\mathcal{A}}^b(\mathcal{A})$ and $D_{\mathcal{A}}^b(\mathcal{P}(A))$), there are shifted dualities, $T_s^n \circ *$ and $T_u^n \circ *$. We work with the unsigned duality $T_u^n \circ *$ until we reach section 6. Note however that $H_i(T_s^n P_{\bullet}^*) = H_i(T_u^n P_{\bullet}^*)$ and so much of what we will say is independent of the chosen duality.

Remark 4.2. We quickly review the situation for the categories \mathcal{A} and \mathcal{R} . First note that both of these categories are exact categories with duality and so the Witt groups are defined as in [QSS].

The functor ι induces duality preserving equivalences

$$\iota : (\mathcal{A}^{\vee}, \tilde{\omega}) \longrightarrow (\mathcal{R}(A), T_u^d \circ *, \varpi), \quad \iota : (\mathcal{A}^{\vee}, -\tilde{\omega}) \longrightarrow (\mathcal{R}(A), T_u^d \circ *, -\varpi)$$

of categories which then yield isomorphisms of the corresponding Witt groups

$$\begin{aligned}
 W(\iota) : W(\mathcal{A}, \vee, \tilde{\varpi}) &\xrightarrow{\sim} W(\mathcal{R}(A), T_u^d \circ *, \varpi), \\
 W(\iota) : W(\mathcal{A}, \vee, -\tilde{\varpi}) &\xrightarrow{\sim} W(\mathcal{R}(A), T_u^d \circ *, -\varpi)
 \end{aligned}$$

Finally, we get to our definitions of the Witt group. Given an exact category \mathcal{E} , its derived category will be denoted by $D^b(\mathcal{E})$. For a subcategory \mathcal{C} , $D_{\mathcal{C}}^b(\mathcal{E})$ will denote the full subcategory of $D^b(\mathcal{E})$ consisting of complexes with homologies in \mathcal{C} .

The derived categories which will be used in this article include:

$$D_{\mathcal{A}}^b(\mathcal{P}(A)) \subseteq D_{fl}^b(\mathcal{P}(A)) \subseteq D^b(\mathcal{P}(A)) \quad D_{\mathcal{A}}^b(\mathcal{A}) \subseteq D^b(\mathcal{A}).$$

We have the following diagram of subcategories and functors:

$$\begin{array}{ccccc}
 \mathcal{A} & \xrightarrow{\mu} & D_{\mathcal{A}}^b(\mathcal{A}) & \xrightarrow{\nu} & D^b(\mathcal{A}) \\
 \downarrow \iota & \searrow \zeta & \downarrow \alpha & & \downarrow \beta \\
 \mathcal{R}(A) & \xrightarrow{\mu'} & D_{\mathcal{A}}^b(\mathcal{P}(A)) & \xrightarrow{\nu'} & D_{fl}^b(\mathcal{P}(A)).
 \end{array} \tag{5}$$

Here $\mu(M)$ is the complex concentrated at degree zero. The functor $\iota(M)$ is obtained by making a choice of projective resolution of length d and then defining $M^\vee = H_0((\zeta(M))^*)$. The functors α and β are essentially induced by these ones, by taking the total complex (take a look at the proof of (3.5)).

We now move on towards the definitions of the Witt groups of the categories $D_{\mathcal{A}}^b(\mathcal{A})$ and $D_{\mathcal{A}}^b(\mathcal{P}(A))$. We once again remind the reader that this definition relies on the definitions in [B2].

Definition 4.3. Let $\delta = \pm 1$. Suppose $K := (K, \#, \delta, \varpi)$ is a triangulated category with translation T and δ -duality $\#$. Suppose K_0 is a full subcategory of K that is closed under isomorphism, translation and orthogonal sum. We abuse notation and denote $K_0 := (K_0, \#, \delta, \varpi)$ in order to keep track of the duality and canonical isomorphism in use.

1. Define the Witt monoid of $MW(K_0)$ to be the submonoid

$$MW(K_0) = \{(P, \varphi) \in MW(K) : P \in Ob(K_0)\}.$$

2. A symmetric space $(P, \varphi) \in MW(K_0)$ will be called a **neutral space** in $MW(K_0)$ if it has a Lagrangian (L, α, w) in $MW(K)$ such that $L, L^\# \in Ob(K_0)$.
3. Let $NW(K_0)$ be the submonoid of $MW(K_0)$ generated by the isometry classes of neutral spaces in K_0 .

4. Define the Witt group

$$W(K_0) := \frac{MW(K_0)}{NW(K_0)}.$$

Note $(Q, \chi) \in MW(K_0) \implies (Q, -\chi) \in MW(K_0)$. It is easy to check that $(Q, \chi) \perp (Q, -\chi) \in NW(K_0)$. So, $W(K_0)$ has a group structure. We use this definition in the context of derived categories of exact categories with duality.

5. Let $(\mathcal{C}^\vee, \varpi)$ be an exact subcategory with duality in an ambient abelian category \mathcal{C}' and let \mathcal{D} be any subcategory of \mathcal{C}' closed under orthogonal sum. Let $K_0 = D_{\mathcal{D}}^b(\mathcal{C})$ (2.1). Then with the induced duality and natural isomorphism, the Witt group $W(D_{\mathcal{D}}^b(\mathcal{C})^\vee, \delta, \varpi)$ is defined as above.
6. Accordingly, with $T = T_u, T_s$, the Witt groups

$$W(D_{\mathcal{A}}^b(\mathcal{P}(A)), T^{n \circ *}, \pm 1, \pm \varpi), \quad W(D_{\mathcal{A}}^b(\mathcal{A}), T^{n \circ *}, \pm 1, \pm \tilde{\varpi})$$

are defined.

5. Isomorphisms of Witt groups

All the functors above induce homomorphisms of Witt groups. As always, **A denotes a Cohen-Macaulay ring with $\dim A_m = d \geq 2$, for all maximal ideals m of A .** Let $D_{\mathcal{A}}^b(\mathcal{A}^\vee, \pm \tilde{\varpi})$, $D^b(\mathcal{A}^\vee, \pm \tilde{\varpi})$ denote the duality structure, respectively, on $D_{\mathcal{A}}^b(\mathcal{A})$ and $D^b(\mathcal{A})$ induced by $(\mathcal{A}^\vee, \pm \tilde{\varpi})$ and

$$D_{\mathcal{A}}^b(\mathcal{P}(A))_u^\pm := (D_{\mathcal{A}}^b(\mathcal{P}(A)), T_u^d \circ *, 1, \pm \varpi).$$

Recall the functors in the diagram (5), it is clear that the functors μ, ν, μ' and γ preserve dualities. For ι , this is left to the reader as a diagram chase (but note the definition of M^\vee after that same diagram). Essentially the same proof also gives us that ζ, α and β are duality preserving. This being done, we can talk about the corresponding maps of the Witt groups. The goal of this section is to establish the following diagram of homomorphisms of Witt groups:

$$\begin{array}{ccccc}
 W(\mathcal{A}^\vee, \pm \tilde{\varpi}) & \xrightarrow[\sim]{W(\mu)} & W(D_{\mathcal{A}}^b(\mathcal{A}^\vee, \pm \tilde{\varpi})) & \xrightarrow[\sim]{W(\nu)} & W(D^b(\mathcal{A}^\vee, \pm \tilde{\varpi})) \\
 W(\iota) \downarrow \wr & \searrow \sim & W(\alpha) \downarrow \wr & & \\
 W(\mathcal{R}(A), T_u^d \circ *, \pm \varpi) & \xrightarrow[\sim]{W(\gamma)} & W(D_{\mathcal{A}}^b(\mathcal{P}(A))_u^\pm) & &
 \end{array} \tag{6}$$

Note that we already know that $W(\iota)$ is an isomorphism (4.2) and further, by [B3, Theorem 4.3], $W(\nu \circ \mu)$ is an isomorphism. The proof that

$$W(\mu) : W(\mathcal{A}^\vee, \pm \tilde{\varpi}) \longrightarrow W(D_{\mathcal{A}}^b(\mathcal{A}^\vee, \pm \tilde{\varpi}))$$

are isomorphisms follows from more abstract “general nonsense” which we prove in (A.1) as part of the appendix (A). Since $W(\nu \circ \mu)$ and $W(\mu)$ are isomorphisms, it is clear that so is $W(\nu)$. The main result of this section is that $W(\zeta)$ is an isomorphism. That being established, it is clear that $W(\gamma)$ and $W(\alpha)$ are isomorphisms.

For the rest of this section, we use the notation $\# := T_u^d \circ *$. First, we establish the following regarding the structure of symmetric forms.

Lemma 5.1. *Suppose $\eta : X_\bullet \xrightarrow{\sim} X_\bullet^\#$ is a symmetric form in $(D_A^b(\mathcal{P}(A))_u^\pm)$, such that*

$$H_{-m}(X_\bullet) \neq 0, \quad \text{and} \quad H_i(X_\bullet) = 0 \quad \text{for all} \quad i < -m.$$

Then, there is a complex P_\bullet in $Ch_A^b(\mathcal{P}(A))$ and a quasi-isomorphism $\varphi : P_\bullet \rightarrow P_\bullet^\#$ such that

1. (P_\bullet, φ) is isometric to (X_\bullet, η) in $(D_A^b(\mathcal{P}(A))_u^\pm)$.
2. P_\bullet is supported on $[m + d, -m]$.
3. $H_{-m}(P_\bullet) \neq 0$.

Proof. Recall from (2.9) that since $H_{-m}(X_\bullet) \neq 0$, X_\bullet has length at least d . By duality, we conclude that $m \geq 0$. By definition there is a complex P_\bullet of projective modules and a quasi-isomorphism $t : P_\bullet \rightarrow X_\bullet$, a chain complex morphism $\varphi_0 : P_\bullet \rightarrow X_\bullet^\#$ such that $\eta = \varphi_0 t^{-1}$. Then, $\varphi = t^\# \varphi_0 = t^\# \eta t$ is a symmetric form on P_\bullet , and (X_\bullet, η) is isometric to (P_\bullet, φ) . By including enough zeros on the two tails, we can assume P_\bullet is supported on $[n + d, -n]$, for some $n \geq m$. If $m = n$ there is nothing to prove. So, assume $n > m$. We have, $H_{-n}(P_\bullet) = 0$. Inductively, we will cut down the support to $[m + d, m]$. We write $\varphi : P_\bullet \rightarrow P_\bullet^\#$ as follows

$$\begin{array}{ccccccccccccccc}
 \longrightarrow & 0 & \longrightarrow & P_{n+d} & \xrightarrow{\partial_{n+d}} & P_{(n-1)+d} & \xrightarrow{\partial_{(n-1)+d}} & \cdots & \xrightarrow{\partial_{-(n-2)}} & P_{-(n-1)} & \xrightarrow{\partial_{-(n-1)}} & P_{-n} & \longrightarrow & 0 & \longrightarrow \\
 & \downarrow & & \downarrow \varphi_n & & \downarrow \varphi_{(n-1)+d} & & & & \downarrow \varphi_{-(n-1)} & & \downarrow \varphi_{-n} & & \downarrow & \\
 \longrightarrow & 0 & \longrightarrow & P_{-n}^* & \xrightarrow{\partial_{-(n-1)}^*} & P_{-(n-1)}^* & \xrightarrow{\partial_{-(n-2)}^*} & \cdots & \xrightarrow{\partial_{(n-1)+d}^*} & P_{(n-1)+d}^* & \xrightarrow{\partial_{n+d}^*} & P_{n+d}^* & \longrightarrow & 0 & \longrightarrow
 \end{array}$$

where P_i are finitely generated projective A -modules. Since $n > m$, $H_{-n}(P_\bullet) \cong H_{-n}(P_\bullet^*) \cong 0$. So, $\partial_{-(n-1)}$ and ∂_{n+d}^* are both split surjections. Thus there are homomorphisms $\epsilon_{-n} : P_{-n} \rightarrow P_{-(n-1)}$ and $\epsilon_{n+d}^* : P_{n+d}^* \rightarrow P_{(n-1)+d}^*$ such that $\partial_{-(n-1)} \circ \epsilon_{-n} = Id$ and $\partial_{n+d}^* \circ \epsilon_{n+d}^* = Id$. Hence, $Z_{-(n-1)}$ and $\frac{P_{n-1+d}}{P_{n+d}}$ are projective modules. Note that since $d \geq 2$, by (2.9), $\frac{P_{n-1+d}}{P_{n+d}} = B_{n-2+d}$. Further, we obtain splittings $\sigma_{-(n-1)} : P_{-(n-1)} \rightarrow Z_{-(n-1)}$ and $\sigma_{(n-2)+d} : B_{(n-2)+d} \hookrightarrow P_{(n-1)+d}$. This gives us a shorter complex Q_\bullet , naturally chain homotopic to P_\bullet and an induced symmetric

form on Q_\bullet :

$$\begin{array}{cccccccccccccccc}
 Q_\bullet : & \rightarrow & 0 & \rightarrow & 0 & \rightarrow & B_{(n-2)+d} & \xrightarrow{c} & P_{(n-2)+d} & \cdots & \xrightarrow{\partial} & Z_{-(n-1)} & \rightarrow & 0 & \rightarrow & 0 & \rightarrow \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow \sigma_{(n-2)+d} & & \parallel & & \downarrow & & \downarrow & & \downarrow & & \downarrow & \\
 P_\bullet : & \rightarrow & 0 & \rightarrow & P_{n+d} & \xrightarrow{\partial_{n+d}} & P_{(n-1)+d} & \xrightarrow{\partial} & P_{(n-2)+d} & \cdots & \xrightarrow{\partial} & P_{-(n-1)} & \xrightarrow{\partial} & P_{-n} & \rightarrow & 0 & \rightarrow \\
 \downarrow \varphi & & \downarrow & & \downarrow \varphi_{n+d} & & \downarrow \varphi_{(n-1)+d} & & \downarrow \varphi_{(n-2)+d} & & \downarrow \varphi_{-(n-1)} & & \downarrow \varphi_{-n} & & \downarrow & & \downarrow & \\
 P_\bullet^\# : & \rightarrow & 0 & \rightarrow & P_{-n}^* & \xrightarrow{\partial_{-(n-1)}^*} & P_{-(n-1)}^* & \xrightarrow{\partial_{-(n-2)}^*} & P_{-(n-2)}^* & \cdots & \xrightarrow{\partial^*} & P_{-(n-1)+d}^* & \xrightarrow{\partial_{n+d}^*} & P_{n+d}^* & \rightarrow & 0 & \rightarrow \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow & & \parallel & & \downarrow \sigma_{(n-2)+d}^* & & \downarrow & & \downarrow & & \downarrow & \\
 Q_\bullet^\# : & \rightarrow & 0 & \rightarrow & 0 & \rightarrow & Z_{-(n-1)}^* & \rightarrow & P_{-(n-2)}^* & \cdots & \xrightarrow{\partial^*} & B_{(n-2)+d}^* & \rightarrow & 0 & \rightarrow & 0 & \rightarrow
 \end{array}$$

Calling this map φ' , (Q_\bullet, φ') is obviously isometric to (P_\bullet, φ) and hence to the original form (X_\bullet, η) . Since Q_\bullet is supported on $[(n-1)+d, -(n-1)]$ induction finishes the proof. \square

Since ζ is given by composing ν and α and there are maps $W(\nu)$ and $W(\alpha)$, it is clear that $W(\zeta)$ is well-defined. However, we give an explicit proof which might also be somewhat illuminating considering the unsaid details about why duality-preserving functors induce maps of Witt groups. The proof essentially follows the proof in [B1, 2.11].

Theorem 5.2. *The functor ζ induces a well defined homomorphism*

$$W(\zeta) : W(\mathcal{A}^\vee, \pm \tilde{\omega}) \longrightarrow W((D_{\mathcal{A}}^b(\mathcal{P}(A))_u^\pm)).$$

Proof. We will only prove

$$W(\zeta) : W(\mathcal{A}^\vee, \tilde{\omega}) \longrightarrow W(D_{\mathcal{A}}^b(\mathcal{A})_u^+).$$

is well defined and the case of skew dualities follows similarly. It is clear that ζ defines a well-defined map from $MW(\mathcal{A}^\vee, \tilde{\omega})$ to $MW(D_{\mathcal{A}}^b(\mathcal{A})_u^+)$ since projective maps of modules can be lifted to a chain complex map of their resolutions (note that though the lift is not unique, it is unique upto homotopy and so gives the same morphism in $D_{\mathcal{A}}^b(\mathcal{P}(A))$). So we need to check that the image of a neutral space in $MW(\mathcal{A}^\vee, \tilde{\omega})$ is neutral in $MW(D_{\mathcal{A}}^b(\mathcal{A})_u^+)$.

Suppose (M, φ_0) is a neutral space in $(\mathcal{A}^\vee, \tilde{\omega})$. Let $\alpha_0 : N \rightarrow M$ be a lagrangian of (M, φ_0) . Then

$$0 \longrightarrow N \xrightarrow{\alpha_0} M \xrightarrow{\alpha_0^\vee \varphi_0} N^\vee \longrightarrow 0 \quad \text{is exact.}$$

Suppose L_\bullet, P_\bullet are the chosen projective resolutions of N and M and $\alpha : L_\bullet \rightarrow P_\bullet$ is the morphism induced from α_0 . The above short exact sequence implies the composition $L_\bullet \xrightarrow{\alpha} P_\bullet \xrightarrow{\alpha^\# \varphi} L_\bullet^\#$ is chain homotopic to 0

(hence the 0 map in $D_{\mathcal{A}}^b(\mathcal{P}(A))$). Completing α to an exact triangle, we get a morphism of exact triangles

$$\begin{array}{ccccccc} L_{\bullet} & \xrightarrow{\alpha} & P_{\bullet} & \xrightarrow{j} & C_{\bullet} & \xrightarrow{k} & T(L_{\bullet}) \\ \downarrow & & \downarrow \alpha^{\#}\varphi & & \downarrow s & & \downarrow \\ 0 & \longrightarrow & L_{\bullet}^{\#} & \xlongequal{\quad} & L_{\bullet}^{\#} & \longrightarrow & 0 \end{array}$$

Note that $H_0(C_{\bullet}) \cong N^{\vee}$ and $\forall i \neq 0 \ H_i(C_{\bullet}) = 0$ and so C_{\bullet} is an object in $D_{\mathcal{A}}^b(\mathcal{P}(A))$. The map s is actually quite easy to describe, namely $s = (0, \alpha^{\#}\varphi) : L_{n-1} \oplus P_n \longrightarrow L_n^*$ and it follows from the above morphism of triangles (or by direct checking) that s is a quasi-isomorphism. Hence,

$$L_{\bullet} \xrightarrow{\alpha} P_{\bullet} \xrightarrow{\alpha^{\#}\varphi} L_{\bullet}^{\#} \xrightarrow{k \circ s^{-1}} T(L_{\bullet})$$

is an exact triangle. Setting $w = -T^{-1}(k \circ s^{-1})$, we get an exact triangle

$$T^{-1}(L_{\bullet}^{\#}) \xrightarrow{w} L_{\bullet} \xrightarrow{\alpha} P_{\bullet} \xrightarrow{\alpha^{\#}\varphi} L_{\bullet}^{\#}$$

Now all we require is that $T^{-1}w^{\#} = w$.

$$\begin{aligned} T^{-1}w^{\#} = w &\Leftrightarrow T^{-1}w^{\#} = -T^{-1}(k \circ s^{-1}) \Leftrightarrow (T^{-1}(k \circ s^{-1}))^{\#} = k \circ s^{-1} \\ &\Leftrightarrow T(s^{-1\#} \circ k^{\#}) = k \circ s^{-1} \Leftrightarrow T(k^{\#}) \circ s = T(s^{\#}) \circ k. \end{aligned}$$

A quick physical check of the maps in question yields that the first map is

$$L_{n-1} \oplus P_n \xrightarrow{\begin{pmatrix} -1 & 0 \end{pmatrix}} L_{n-1} \xrightarrow{\begin{pmatrix} 0 \\ \varphi_{n-1}^* \circ \alpha_{d-n+1} \end{pmatrix}} L_{d-n}^* \oplus P_{d-n+1}^*$$

while the second one is

$$L_{n-1} \oplus P_n \xrightarrow{\begin{pmatrix} 0 & \alpha_{d-n}^* \circ \varphi_n \end{pmatrix}} L_{d-n}^* \xrightarrow{\begin{pmatrix} -1 \\ 0 \end{pmatrix}} L_{d-n}^* \oplus P_{d-n+1}^*$$

The matrices we thus obtain are

$$\begin{pmatrix} 0 & -\varphi_{n-1}^* \circ \alpha_{d-n+1} \\ 0 & 0 \end{pmatrix} \qquad \begin{pmatrix} 0 & 0 \\ \alpha_{d-n}^* \circ \varphi_n & 0 \end{pmatrix}$$

which are homotopy equivalent using

$$\tau = \begin{pmatrix} 0 & 0 \\ 0 & (-1)^n \varphi \end{pmatrix}.$$

Therefore, (L_{\bullet}, α, w) is a lagrangian. Hence $W(\zeta)$ is a well defined homomorphism of groups. □

We now proceed towards (5.11) which proves that $W(\zeta)$ is surjective. The main tool here is to construct a special sublagrangian and then use Balmer’s sublagrangian construction [B2, Section 4 and Theorem 4.20] to reduce the length of (P, φ) .

Remark 5.3. Note that using (5.1), any symmetric form (X_\bullet, ϕ) in $(D_{\mathcal{A}}^b(\mathcal{A})_u^+)$ with X_\bullet not acyclic can be represented by

$$\begin{array}{ccccccccccc}
 P_\bullet = \cdots & 0 & \longrightarrow & P_{n+d} & \xrightarrow{\partial} & P_{(n-1)+d} & \xrightarrow{\partial} & \cdots & \xrightarrow{\partial} & P_{-(n-1)} & \xrightarrow{\partial} & P_{-n} & \longrightarrow & 0 \\
 & \varphi \downarrow & & \varphi_n \downarrow & & \varphi_{(n-1)+d} \downarrow & & & & \varphi_{-(n-1)} \downarrow & & \varphi_{-n} \downarrow & & \\
 P_\bullet^\# = \cdots & 0 & \longrightarrow & P_{-n}^* & \xrightarrow{\partial^*} & P_{-(n-1)}^* & \longrightarrow & \cdots & \xrightarrow{\partial^*} & P_{(n-1)+d}^* & \xrightarrow{\partial^*} & P_{n+d}^* & \longrightarrow & 0
 \end{array}$$

with $H_{-n}(P_\bullet) \neq 0$.

Lemma 5.4. *Let (P_\bullet, φ) be as above. Then*

1. $H_r(P_\bullet) = 0$ for $r = n + 1, n + 2, \dots, n + d$.
2. $H_n(P_\bullet) \neq 0$.

Proof. The first point follows from (2.9). To prove (2), assume $H_n(P_\bullet) = 0$. Then, with $B_{n-1} = \text{image}(\partial_n)$ we have an exact sequence

$$0 \longrightarrow P_{n+d} \longrightarrow P_{(n-1)+d} \longrightarrow \cdots \longrightarrow P_n \longrightarrow P_{n-1} \longrightarrow \frac{P_{n-1}}{B_{n-1}} \longrightarrow 0$$

Since this is a projective resolution of the last term, it follows

$$H_{-n}(P_\bullet^*) = \text{Ext}^{d+1} \left(\frac{P_{n-1}}{B_{n-1}}, A \right) = 0.$$

This is a contradiction to $H_{-n}(P_\bullet) \neq 0$. The proof is complete. □

Much of what follows is dependent on [B2, §4] and the interested reader is highly encouraged to take a look at it. We recall the definition of a sublagrangian of (P_\bullet, φ) :

Definition 5.5. *A sublagrangian of a symmetric form (P_\bullet, φ) is a pair (L_\bullet, α) with $L_\bullet \in \text{Ob}(D_{\mathcal{A}}^b(\mathcal{P}(A)))$ and $\alpha : L_\bullet \rightarrow P_\bullet$, which satisfies that $\alpha^\# \circ \varphi \circ \alpha = 0$ in $D_{\mathcal{A}}^b(\mathcal{P}(A))$.*

For (P_\bullet, φ) as above, (5.4) tells us that $H_n(P_\bullet) \neq 0$ and we already know it is in \mathcal{A} . So it has a minimal projective resolution of length d . Let L_\bullet be a projective resolution of $H_n(P_\bullet)$ of length d , shifted by n places, as in the diagram below. Since $H_i(P_\bullet) \forall i > n$ by (2.9), the bottom line is a

Proof. (1) follows directly from (3.1). (2) follows because by (5.7) N_\bullet is supported on $[(n+1)+d, -n]$. For (3), notice that the only nonzero homology of $L_\bullet^\#$ is at degree $-n$. Since $n > 0$, the long exact homology sequence of the second triangle gives us

$$H(v_2^\#) : H_n(N_\bullet^\#) \xrightarrow{\sim} H_n(P_\bullet^\#).$$

By choice of v and φ , we know that

$$H(v) : H_n(L_\bullet) \xrightarrow{\sim} H_n(P_\bullet), \quad H(\varphi) : H_n(P_\bullet) \xrightarrow{\sim} H_n(P_\bullet^\#)$$

and hence, the commutative diagram

$$\begin{array}{ccc} H_n(L_\bullet) & \xrightarrow[\sim]{H_n(v)} & H_n(P_\bullet) \\ \downarrow H_n(\mu_0) & & \downarrow H_n(\varphi) \\ H_n(N_\bullet^\#) & \xrightarrow[\sim]{H_n(v_2^\#)} & H_n(P_\bullet^\#) \end{array}$$

gives us (3). We prove (4) now. Since the only nonzero homology of $L_\bullet^\#$ is at degree $-n$, it is clear from the long exact homology sequence for the bottom exact triangle that

$$H_i(N_\bullet^\#) \cong H_i(P_\bullet^\#) \quad \forall i \neq -n, -n - 1.$$

By (5.7), $H_i(N_\bullet) = 0$ for all $i \geq n$ and so

$$H_{-(n+1)}(N_\bullet^\#) = \text{Ext}^{d+2} \left(\frac{N_{n-1}}{B_{n-1}}, A \right) = 0, \quad H_{-n}(N_\bullet^\#) = \text{Ext}^{d+1} \left(\frac{N_{n-1}}{B_{n-1}}, A \right) = 0.$$

where $B_{n-1} \subseteq N_{n-1}$ is the boundary submodule (the last part also follows directly because $\text{Ext}^d(\frac{P_n}{B_n}, A) \cong \text{Ext}^d(H_n(P_\bullet), A)$). So, (4) is established. The proof is complete. \square

Now we consider the cone of μ_0 .

Lemma 5.9. *With the notations in (5.6), (5.7) and (5.8), consider an exact triangle on μ_0 as follows:*

$$L_\bullet \xrightarrow{\mu_0} N_\bullet^\# \xrightarrow{\mu_1} R_\bullet \xrightarrow{\mu_2} T(L_\bullet)$$

where R_\bullet is the cone of μ_0 . Then R_\bullet is an object of $D_{\mathcal{A}}^b(\mathcal{P}(A))$. More precisely,

$$H_i(R_\bullet) = \begin{cases} H_i(N_\bullet^\#) & \text{for } -(n-1) \leq i \leq n-1 \\ 0 & \text{otherwise.} \end{cases}$$

which tells us that R_\bullet has exactly two nonzero homologies less than than P_\bullet .

Now, $H_i(R_\bullet) = 0$ if $i < -(n - 1), i > (n - 1)$. By (5.1), $(R_\bullet, -\psi)$ is isometric to a form (Q_\bullet, τ) such that Q_\bullet is supported on $[k + d, -k]$ with $0 \leq k \leq n - 1$. The proof is complete. \square

Now we are ready to state and prove the main result of this article, which is our version of the dévissage theorem, i.e.

Theorem 5.12. *The homomorphisms*

$$W(\zeta) : W(\mathcal{A}^\vee, \pm\tilde{\omega}) \longrightarrow W((D_{\mathcal{A}}^b(\mathcal{P}(A))_u^\pm))$$

induced by the functor ζ are isomorphisms.

Proof. First, we prove the surjectivity of the homomorphism $W(\zeta)$. Suppose $x \in W((D_{\mathcal{A}}^b(\mathcal{P}(A))_u^\pm))$. Then, by (5.1), we can write $x = [(P_\bullet, \varphi)]$ of the form described in (5.3). Inductively, by (5.11), there is a form (R_\bullet, ψ) in $(D_{\mathcal{A}}^b(\mathcal{P}(A))_u^\pm)$ such that

$$[(R_\bullet, \psi)] = [(P_\bullet, \varphi)] = x \quad \text{in} \quad W((D_{\mathcal{A}}^b(\mathcal{P}(A))_u^\pm))$$

and R_\bullet is supported in $[d, 0]$. By (2.9), R_\bullet is a projective resolution of $M := H_0(R_\bullet) \in \mathcal{A}$. Further, ψ induces a form $\psi_0 : M \xrightarrow{\sim} M^\vee$ and clearly

$$W(\zeta)([(M, \psi_0)]) = [(R_\bullet, \psi)] = x.$$

So, $W(\zeta)$ is surjective.

Now, we proceed to prove that $W(\zeta)$ is injective. Suppose (M, q) is a symmetric form in $(\mathcal{A}^\vee, \pm\tilde{\omega})$ and $W(\zeta)([(M, q)]) = 0$. Write, $(\zeta(M), \zeta(q)) = (P_\bullet, \varphi_0)$ where P_\bullet is a projective resolution of M of length d , and φ_0 is the induced symmetric form. So, $[(P_\bullet, \varphi_0)] = 0$ in $W((D_{\mathcal{A}}^b(\mathcal{P}(A))_u^\pm))$.

This means there is a neutral form $[(Q_\bullet, \varphi_1)]$ so that $[(P_\bullet, \varphi_0)] \perp [(Q_\bullet, \varphi_1)]$ is neutral in $W((D_{\mathcal{A}}^b(\mathcal{P}(A))_u^\pm))$. Since $[(Q_\bullet, \varphi_1)]$ is neutral, so is $[(Q_\bullet, -\varphi_1)]$. Hence, $[(P_\bullet, \varphi_0)] \perp [(Q_\bullet, \varphi_1)] \perp [(Q_\bullet, -\varphi_1)]$ is neutral. Using the usual isometry, we get that there is a hyperbolic form

$$\left(Q_\bullet \oplus Q_\bullet^\#, \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right) \quad \text{with} \quad Q_\bullet \in D_{\mathcal{A}}^b(\mathcal{P}(A))$$

such that

$$(U_\bullet, \varphi) := \left(P_\bullet \oplus Q_\bullet \oplus Q_\bullet^\#, \begin{pmatrix} \varphi_0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \right)$$

is neutral in $W((D_{\mathcal{A}}^b(\mathcal{P}(A))_u^\pm))$

(we have so far followed the argument in [B2, 3.5]).

So, (U_\bullet, φ) has a lagrangian (L_\bullet, α) . We have an exact triangle

$$T^{-1}L_\bullet \xrightarrow{w} L_\bullet \xrightarrow{\alpha} U_\bullet \xrightarrow{\alpha^\# \varphi} L_\bullet \quad T^{-1}w^\# = w.$$

Before proceeding, we use (3.3) to make a startlingly simple observation about the homologies of $L_\bullet^\#$:

$$H_i(L_\bullet^\#) \cong H_{i-d}(L_\bullet^*) \cong \text{Ext}^d(H_{(d-i)-d}(L_\bullet), A) \cong \text{Ext}^d(H_{-i}(L_\bullet), A) \cong H_{-i}(L_\bullet)^\vee.$$

Similarly, $H_i(U_\bullet^\#) \cong H_{-i}(U_\bullet)^\vee$ and further using the remark (3.4) about naturality of the maps, we get that the long exact homology sequences are

$$\begin{array}{ccccccc} \longrightarrow & H_{-2}(L_\bullet)^\vee & \xrightarrow{H_1(w)} & H_1(L_\bullet) & \xrightarrow{H_1(\alpha)} & H_1(U_\bullet) & \xrightarrow{H_{-1}(\alpha)^\vee \circ H_1(\varphi)} & H_{-1}(L_\bullet)^\vee & \cdots \\ & \parallel & & \tilde{\varphi}_{H_1} \downarrow \wr & & H_1(\varphi) \downarrow \wr & & \parallel & \\ \longrightarrow & H_{-2}(L_\bullet)^\vee & \xrightarrow{H_{-2}(w)^\vee} & H_1(L_\bullet)^{\vee\vee} & \xrightarrow{H_{-1}(\varphi)^\vee \circ H_1(\alpha)^\vee} & H_{-1}(U_\bullet)^\vee & \xrightarrow{H_{-1}(\alpha)^\vee} & H_{-1}(L_\bullet)^\vee & \cdots \\ & & & & & & & & \\ \cdots & \xrightarrow{H_0(w)} & H_0(L_\bullet) & \xrightarrow{H_0(\alpha)} & H_0(U_\bullet) & \xrightarrow{H_0(\alpha)^\vee \circ H_0(\varphi)} & H_0(L_\bullet)^\vee & \xrightarrow{H_{-1}(w)} & \cdots \\ & & \tilde{\varphi}_{H_0} \downarrow \wr & & H_0(\varphi) \downarrow \wr & & \parallel & & \\ \cdots & \xrightarrow{H_{-1}(w)^\vee} & H_0(L_\bullet)^{\vee\vee} & \xrightarrow{H_0(\varphi)^\vee \circ H_0(\alpha)^\vee} & H_0(U_\bullet)^\vee & \xrightarrow{H_0(\alpha)^\vee} & H_0(L_\bullet)^\vee & \xrightarrow{H_0(w)^\vee} & \cdots \\ & & & & & & & & \\ \cdots & H_{-1}(L_\bullet) & \xrightarrow{H_{-1}(\alpha)} & H_{-1}(U_\bullet) & \xrightarrow{H_1(\alpha)^\vee \circ H_{-1}(\varphi)} & H_{-1}(L_\bullet)^\vee & \xrightarrow{H_{-2}(w)} & H_{-2}(L_\bullet) & \xrightarrow{H_{-2}(\alpha)} & \cdots \\ & \tilde{\varphi}_{H_{-1}} \downarrow \wr & & H_{-1}(\varphi) \downarrow \wr & & \parallel & & \tilde{\varphi}_{H_{-2}} \downarrow \wr & & \\ \cdots & H_{-1}(L_\bullet)^{\vee\vee} & \xrightarrow{H_{-1}(\varphi)^\vee \circ H_{-1}(\alpha)^\vee} & H_1(U_\bullet)^\vee & \xrightarrow{H_1(\alpha)^\vee} & H_1(L_\bullet)^\vee & \xrightarrow{H_1(w)^\vee} & H_{-2}(L_\bullet)^{\vee\vee} & \xrightarrow{H_2(\varphi)^\vee \circ H_{-2}(\alpha)^\vee} & \cdots \end{array}$$

Replacing the part of the top exact sequence in negative degree by the corresponding part of the bottom (dual) exact sequence, we get an exact sequence:

$$\begin{array}{ccccccc} \longrightarrow & H_{-2}(L_\bullet)^\vee & \xrightarrow{H_1(w)} & H_1(L_\bullet) & \xrightarrow{H_1(\alpha)} & H_1(U_\bullet) & \xrightarrow{H_{-1}(\alpha)^\vee \circ H_1(\varphi)} & H_{-1}(L_\bullet)^\vee \\ & & & & & & & \\ \xrightarrow{H_0(w)} & H_0(L_\bullet) & \xrightarrow{H_0(\alpha)} & H_0(U_\bullet) & \xrightarrow{H_0(\alpha)^\vee \circ H_0(\varphi)} & H_0(L_\bullet)^\vee & \xrightarrow{H_0(w)^\vee} & H_{-1}(L_\bullet)^{\vee\vee} \\ & & & & & & & \\ \xrightarrow{H_1(\varphi)^\vee \circ H_{-1}(\alpha)^\vee} & H_1(U_\bullet)^\vee & \xrightarrow{H_1(\alpha)^\vee} & H_1(L_\bullet)^\vee & \xrightarrow{H_1(w)^\vee} & H_{-2}(L_\bullet)^{\vee\vee} & \xrightarrow{H_2(\varphi)^\vee \circ H_{-2}(\alpha)^\vee} & \cdots \end{array}$$

Notice that the complex above is very special, and is “symmetric” about $H_0(U_\bullet)$. So we can apply [B3, 4.1 Lemma] to this sequence. Since the sequence is exact, we have

$$[(H_0(U_\bullet), H_0(\varphi))] = [(0, 0)] = 0 \quad \text{in } W(\mathcal{MFPD}^f(A)).$$

However,

$$\begin{aligned} (H_0(U_\bullet), H_0(\varphi)) &= \left(H_0(P_\bullet) \oplus H_0(Q_\bullet) \oplus H_0(Q_\bullet^\#), \begin{pmatrix} H_0(\varphi_0) & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \right) \\ &= (M, q) \perp \left(H_0(Q_\bullet) \oplus H_0(Q_\bullet)^\vee, \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right) \end{aligned}$$

So, we have

$$\begin{aligned} [(M, q)] &= \left[(M, q) \perp \left(H_0(Q_\bullet) \oplus H_0(Q_\bullet)^\vee, \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right) \right] \\ &= [(H_0(U_\bullet), H_0(\varphi))] = 0. \end{aligned}$$

The proof is complete. □

6. Shifted Witt groups

In this section, we use the previous results to obtain our dévissage theorem for the Witt groups $W^i(D_{\mathcal{A}}^b(\mathcal{P}(A)))$. We recall that A is a Cohen-Macaulay ring with $\dim A_m = d \geq 2$ for all maximal ideals m and such that 2 is invertible in A and that $\mathcal{A} = \mathcal{MFPD}^{fl}(A)$.

Notations 6.1. For integers $j \geq 0$ define the functor $\zeta_j = T^{-j} \circ \zeta$, which associates to an object M in \mathcal{A} a projective resolution P_\bullet of M of length d , such that $H_{-j}(P_\bullet) = M$.

Definition 6.2. Suppose $K := (K, \#, \delta, \varpi)$ is a triangulated category with translation T and δ -duality $\#$. We recall from [B3] that

$$T^n K := \left(K, T^n \circ \#, (-1)^n \delta, (-1)^{\frac{n(n+1)}{2}} \delta^n \varpi \right).$$

is then also a triangulated category with the same translation T but with $((-1)^n \delta)$ -duality $T^n \circ \#$. If K_0 is a subcategory of K satisfying the conditions of (4.3), we define $T^n K_0$ to be the same subcategory and translation with the induced duality structure from $T^n K$. Using (4.3), we define the shifted Witt groups by

$$W^n(K) := W(T^n K) \quad W^n(K_0) := W(T^n K_0) \quad \forall n \in \mathbb{Z}.$$

Note that $T_s^2 : T^n K \rightarrow T^{n+4} K$ is an equivalence of triangulated categories with duality, for all $n \in \mathbb{Z}$. Similarly, $T_s^2 : T^n K_0 \rightarrow T^{n+4} K_0$ is an equivalence of categories with duality, for all $n \in \mathbb{Z}$ and so

$$W^n(K) \xrightarrow{\sim} W^{n+4}(K) \quad W^n(K_0) \xrightarrow{\sim} W^{n+4}(K_0) \quad \forall n \in \mathbb{Z}.$$

Definition 6.3. Following [BW], by “**standard**” duality structure on \mathcal{A} , we mean the exact category $(\mathcal{A}^\vee, (-1)^{\frac{d(d-1)}{2}} \tilde{\omega})$. By “**standard**” skew duality structure on \mathcal{A} , we mean the exact category $(\mathcal{A}^\vee, -(-1)^{\frac{d(d-1)}{2}} \tilde{\omega})$. We denote the Witt groups

$$W_{St}^+(A) = W(\mathcal{A}^\vee, (-1)^{\frac{d(d-1)}{2}} \tilde{\omega}), \quad W_{St}^-(A) = W(\mathcal{A}^\vee, -(-1)^{\frac{d(d-1)}{2}} \tilde{\omega}).$$

Theorem 6.4. *Then, the functor $\zeta_0 : \mathcal{A} \rightarrow D_{\mathcal{A}}^b(\mathcal{P}(A))$ induces an isomorphism*

$$W(\zeta_0) : W_{St}^+(A) \xrightarrow{\sim} W^d(D_{\mathcal{A}}^b(\mathcal{P}(A)), *, 1, \varpi).$$

Proof. Recall ζ_0 was denoted by ζ in the previous sections. For notational convenience $\varpi_0 = (-1)^{\frac{d(d-1)}{2}} \varpi$. By theorem (5.12), we get the following isomorphism of Witt groups

$$\eta_0 : W_{St}^+(\mathcal{A}) \xrightarrow{\sim} W(D_{\mathcal{A}}^b(\mathcal{P}(A)), \#_d^\mu, 1, \varpi_0).$$

There is a duality preserving equivalence [BW, Proof of Lemma 6.4]

$$\beta : (D^b(\mathcal{P}(A)), T_u^d \circ *, 1, \varpi_0) \rightarrow (D^b(\mathcal{P}(A)), T_u^d \circ *, (-1)^d, (-1)^{\frac{d(d+1)}{2}} \varpi).$$

Note that the later is the shifted category $T^d(D^b(\mathcal{P}(A)), *, 1, \varpi)$. Composing η_0 with the homomorphism induced by β , we get the result. \square

Now we prove the standard skew duality version of theorem (6.4).

Theorem 6.5. *The functor ζ_1 induces an isomorphism*

$$W_{St}^-(A) \xrightarrow{\sim} W^{d-2}(D_{\mathcal{A}}^b(\mathcal{P}(A)), *, 1, -\varpi).$$

Proof. By Theorem (5.12), we have an isomorphism

$$W_{St}^-(A) \xrightarrow{\sim} W(D_{\mathcal{A}}^b(\mathcal{P}(A)), T_u^d \circ *, 1, -(-1)^{\frac{d(d-1)}{2}} \varpi).$$

Write $\varpi_0 = -(-1)^{\frac{d(d-1)}{2}} \varpi$. There is an equivalence of categories [B2, 2.14]

$$T_s : (D_{\mathcal{A}}^b(\mathcal{P}(A)), T_u^{d-2} \circ *, 1, \varpi_0) \rightarrow (D_{\mathcal{A}}^b(\mathcal{P}(A)), T_u^d \circ *, 1, \varpi_0).$$

This induces an isomorphism

$$W(D_{\mathcal{A}}^b(\mathcal{P}(A)), T_u^{d-2} \circ *, 1, \varpi_0) \xrightarrow{\sim} W(D_{\mathcal{A}}^b(\mathcal{P}(A)), T_u^d \circ *, 1, \varpi_0).$$

As in the proof of [BW, Lemma 6.4], there is an equivalence of triangulated categories with duality

$$(D^b(\mathcal{P}(A)), T_u^{d-2} \circ *, 1, \varpi_0) \rightarrow (D^b(\mathcal{P}(A)), T_s^{d-2} \circ *, (-1)^{d-2}, (-1)^{\frac{d(d+1)}{2}} \varpi).$$

The latter category is $T^{d-2}(D^b(\mathcal{P}(A)), *, 1, -\varpi)$. The proof is complete. \square

Finally, we have the following regarding odd shifts.

Theorem 6.6. *For $n = d - 1, d - 3$, we have*

$$W^n(D_{\mathcal{A}}^b(\mathcal{P}(A)), *, 1, \pm\varpi) = 0.$$

Proof. First consider $n = d - 1$. It would be enough to prove that

$$W(D_{\mathcal{A}}^b(\mathcal{P}(A)), T_u^{d-1} \circ *, 1, \pm\varpi) = 0.$$

Suppose (P_{\bullet}, φ) is a form in $D_{\mathcal{A}}^b(\mathcal{P}(A), T_u^{d-1} \circ *, 1, \pm\varpi)$. By a little tweak in (5.1), we can assume that P_{\bullet} is supported on $[n + (d - 1), -n]$ with $n > 0$ and $H_{-n}(P_{\bullet}) \neq 0$. By imitating the arguments of theorem (5.11), we can keep shortening the length of the complexes which give our symmetric form. Eventually, we will be reduced to the case where the complex is P_{\bullet} is supported on $[d - 1, 0]$. By theorem (2.9), P_{\bullet} is exact. So, $[(P, \varphi)] = 0$. The same arguments apply when $n = d - 3$. The proof is complete. \square

Using the 4-periodicity, we now obtain the theorem mentioned in the introduction:

Theorem 6.7 (shiftFinal). *Let $\mathcal{B} = (D_{\mathcal{A}}^b(\mathcal{P}(A)), T_s, *, 1, \varpi)$. Then, for $n \in \mathbb{Z}$, we have*

1. $W^{d+4n}(\mathcal{B}) = W_s^+(\mathcal{A})$,
2. $W^{d+4n+1}(\mathcal{B}) = 0$,
3. $W^{d+4n+2}(\mathcal{B}) = W_s^-(\mathcal{A})$,
4. $W^{d+4n+3}(\mathcal{B}) = 0$.

A. Some formalism

The purpose of this section is to prove the following theorem:

Theorem A.1. *Suppose \mathcal{E} is a full subcategory of a $\mathbb{Z}[\frac{1}{2}]$ abelian category \mathcal{B} with the 2 out of 3 property for short exact sequences, and has duality $(\mathcal{E}, {}^\vee, \tilde{\varpi})$. Let $D^b(\mathcal{E}) := (D^b(\mathcal{E}), *, a, \varpi)$ denote the derived category, with duality, of $(\mathcal{E}, {}^\vee, \tilde{\varpi})$. Also, let $D_{\mathcal{E}}^b(\mathcal{E})$ denote the derived category, with duality, of objects in $D^b(\mathcal{E})$ with homologies in \mathcal{E} . Then the homomorphism*

$$W(\mu) : W(\mathcal{E}, {}^\vee, \tilde{\varpi}) \longrightarrow W(D_{\mathcal{E}}^b(\mathcal{E}))$$

induced by the functor $\mu : \mathcal{E} \longrightarrow D_{\mathcal{E}}^b(\mathcal{E})$ is an isomorphism.

In particular, with $\mathcal{E} = \mathcal{A}$ and $\mathcal{B} = \mathcal{M}(A)$, we obtain that

$$W(\mu) : W(\mathcal{A}, {}^\vee, \pm\tilde{\varpi}) \longrightarrow W(D_{\mathcal{A}}^b(\mathcal{A}, {}^\vee, \pm\tilde{\varpi}))$$

as promised in section 5.

The proof of the theorem is essentially the same as the proof of [B3, Theorem 3.2] with the extra check that all constructions yield complexes whose homologies are in \mathcal{A} . This boils down to using the most elementary of sublagrangians (concentrated in just one degree) and reducing length. In any case, we follow the proof in [B3, Theorem 3.2]. Since the category \mathcal{E} has all the properties required in the results in [B3, Section 3]), we will freely borrow them.

To start with, injectivity of $W(\mu)$ follows directly because the isomorphism $W(\mathcal{E}) \xrightarrow{\sim} W(D^b(\mathcal{E}))$ (proven in [B3, Theorem 4.3]) factors as

$$\begin{array}{ccc} W(\mathcal{E}) & \xrightarrow{W(\mu)} & W(D_{\mathcal{E}}^b(\mathcal{E})) \\ & \searrow \sim & \downarrow \\ & & W(D^b(\mathcal{E})) \end{array}$$

We move to the proof of surjectivity which, as we mentioned above will require checking that we remain in $D_{\mathcal{E}}^b(\mathcal{E})$ through all the lemmas establishing [B3, Theorem 3.2]. To start with, we establish the following result regarding duality, which also provides an alternative proof of (3.5).

Lemma A.2. *With the same notations as in (A.1), $D_{\mathcal{E}}^b(\mathcal{E})$ is closed under duality.*

Proof. Let P_{\bullet} be an object in the derived category $D_{\mathcal{E}}^b(\mathcal{E})$. Write

$$P_{\bullet} : \quad \cdots \longrightarrow P_2 \xrightarrow{d_2} P_1 \xrightarrow{d_1} P_0 \longrightarrow P_{-1} \longrightarrow P_{-2} \longrightarrow \cdots$$

$$P_{\bullet}^* : \quad \cdots \longrightarrow P_{-2}^{\vee} \longrightarrow P_{-1}^{\vee} \longrightarrow P_0^{\vee} \longrightarrow P_1^{\vee} \longrightarrow P_2^{\vee} \longrightarrow \cdots$$

Since the complexes are bounded and the homologies are objects of \mathcal{E} , all the kernels $Z_i = \ker(d_i)$, images $B_i = \text{image}(d_{i+1})$ and quotients $\frac{P_i}{B_i}$ are also objects of \mathcal{E} . Hence, so are their duals. But we have an exact sequence

$$0 \longrightarrow \left(\frac{P_{t-1}}{B_{t-1}}\right)^{\vee} \longrightarrow P_{t-1}^{\vee} \longrightarrow \left(\frac{P_t}{B_t}\right)^{\vee} \longrightarrow H_t(P_{\bullet}^*) \longrightarrow 0$$

The first three terms in this sequence are in \mathcal{E} , hence so is $H_0(P_{\bullet}^*)$. The proof is complete. □

Lemma A.3. *Let $x \in W(D_{\mathcal{E}}^b(\mathcal{E}))$. Then $x = (P_{\bullet}, s)$ such that*

1. P_{\bullet} is bounded and $s : P_{\bullet} \longrightarrow P_{\bullet}^*$ is a morphism of complexes, without denominator.

2. s is quasi-isomorphism.
3. s is strongly symmetric (i.e $s_{-i}^\vee = s_i \forall i \in \mathbb{Z}$).
4. $H_i(P_\bullet) \in \mathcal{E}$ for all $i \in \mathbb{Z}$.

Proof. Let the form x be given by (X_\bullet, η) . By definition there is a complex P_\bullet which is an object of $D_{\mathcal{E}}^b(\mathcal{E})$ and a chain complex quasi-isomorphisms $t : P_\bullet \rightarrow X_\bullet$ and $\varphi_0 : P_\bullet \rightarrow X_\bullet^*$ such that $\eta = \varphi_0 t^{-1}$. Then, $s = t^* \varphi_0 = t^* \eta t$ is a symmetric form on P_\bullet and (X_\bullet, η) is isometric to (P_\bullet, φ) . Clearly s is an actual morphism of complexes, a quasi-isomorphism and $H_i(P_\bullet) \in \mathcal{E}$ for all $i \in \mathbb{Z}$. Finally, using that $\frac{1}{2}$ exists, we can make the map strongly symmetric. \square

Lemma A.4. *Let (P_\bullet, s) be a symmetric form in $D_{\mathcal{E}}^b(\mathcal{E})$ as in (A.3), such that P_\bullet is supported on $[m, -n]$ with $m > n \geq 0$. Then (P_\bullet, s) is isometric to a symmetric space (Q_\bullet, t) such that Q_\bullet is supported on $[n, -n]$ and (Q_\bullet, t) has all the other properties of (P_\bullet, s) .*

Proof. This is precisely [B3, Lemma 3.7] in our context. Note that since we can do this in the derived category without the homology condition $D^b(\mathcal{E})$, we use the same result to get (Q_\bullet, t) isometric to (P_\bullet, s) with the required condition. However, since the isometry gives in particular a quasi-isomorphism $P_\bullet \xrightarrow{\sim} Q_\bullet$ and so Q_\bullet is also an object in $D_{\mathcal{E}}^b(\mathcal{E})$. The proof is complete. \square

Lemma A.5. *Let (P_\bullet, s) be a symmetric space, as in (A.3). with support on $[-n, n]$ and $n > 0$. Then there exists a symmetric space (Q_\bullet, t) such that*

1. (Q_\bullet, t) is as in (A.3).
2. (Q_\bullet, t) is supported in $[n, -(n - 1)]$.
3. $H_i(Q_\bullet) \in \mathcal{E}$ for all $i \in \mathbb{Z}$.
4. $[(P_\bullet, s)] + [(Q_\bullet, t)] = 0$ in $W(D_{\mathcal{E}}^b(\mathcal{E}))$.

Proof. Once again this is [B3, Lemma 3.9] in our context. We skim through the proof mentioning only the significant points and most important, checking the points where we need to check the extra homology condition. We begin by proving the lemma in the case $n \geq 2$. Write

$$\begin{array}{ccccccccccc}
 P_\bullet = & & \dots & 0 & \longrightarrow & P_n & \xrightarrow{d} & P_{n-1} & \xrightarrow{d} & \dots & \xrightarrow{d} & P_{-n} & \longrightarrow & 0 \\
 \downarrow s & & & & & \downarrow s & & \downarrow s & & & & \downarrow s & & & \\
 P_\bullet^* = & & \dots & 0 & \longrightarrow & P_{-n}^\vee & \xrightarrow{d^\vee} & P_{-(n-1)}^\vee & \xrightarrow{d^\vee} & \dots & \longrightarrow & P_n^\vee & \longrightarrow & 0
 \end{array}$$

Define (Q_\bullet, t) as follows, on the left side:

$$\begin{array}{ccccccccccccccc}
 Q_\bullet = \cdots & 0 & \longrightarrow & P_n & \xrightarrow{\begin{pmatrix} s \\ d \end{pmatrix}} & P_{-n}^V \oplus P_{n-1} & \xrightarrow{(0,d)} & P_{n-2} & \xrightarrow{d} & \cdots & P_{-(n-2)} & \xrightarrow{d} & P_{-(n-1)} & \longrightarrow & 0 & \longrightarrow & 0 \\
 \downarrow t & & & \downarrow 0 & & \downarrow (d^V, -s) & & \downarrow -s & & & \downarrow -s & & \downarrow \begin{pmatrix} d \\ -s \end{pmatrix} & & & & & \\
 Q_\bullet^* = \cdots & 0 & \longrightarrow & 0 & \xrightarrow{d^V} & P_{-(n-1)}^V & \xrightarrow{d^V} & P_{-(n-2)}^V & \xrightarrow{d^V} & \cdots & P_{(n-2)}^V & \xrightarrow{\begin{pmatrix} 0 \\ d^V \end{pmatrix}} & P_{-n} \oplus P_{n-1}^V & \xrightarrow{P_{n-1}^V} & P_n^V & \longrightarrow & 0
 \end{array}$$

It was proved in [B3, Lemma 3.9] that t is a quasi-isomorphism. So, it follows

$$H_n(Q_\bullet) \cong 0, \quad H_{n-1}(Q_\bullet) \cong H_{n-1}(Q_\bullet^*) \cong \ker(d^V) \in \mathcal{E}.$$

Since $\text{image}(0, d_{n-1}) = \text{image}(d_{n-1})$ we have

$$H_i(Q_\bullet) = H_i(P_\bullet) \in \mathcal{E} \quad \forall \quad i \leq n - 2.$$

Therefore

$$H_i(Q_\bullet) \in \mathcal{E} \quad \text{for all } i \in \mathbb{Z}.$$

So Q_\bullet satisfies the last condition of (A.3). The other conditions of (A.3) are shown to be established in [B3, Lemma 3.9].

Therefore, Q_\bullet satisfies (A.3). It was established in [B3] that $(P_\bullet, s) \perp (Q_\bullet, t)$ is neutral in $D^b(\mathcal{E})$, by showing that $(P_\bullet, s) \perp (Q_\bullet, t)$ is isometric (in $D^b(\mathcal{E})$) to the cone of the morphism $z : T^{-1}M_\bullet^* \rightarrow M_\bullet$ defined as follows:

$$\begin{array}{ccccccccccccccc}
 T^{-1}M_\bullet^* = \cdots & 0 & \longrightarrow & P_n & \xrightarrow{-d} & P_{n-1} & \xrightarrow{-d} & \cdots & P_{-(n-2)} & \xrightarrow{-d} & P_{-(n-1)} & \longrightarrow & 0 \\
 \downarrow z & & & \downarrow 0 & & \downarrow 0 & & & \downarrow 0 & & \downarrow s_{-n}d & & & \\
 M_\bullet = \cdots & 0 & \longrightarrow & P_{-(n-1)}^V & \xrightarrow{d^V} & P_{-(n-2)}^V & \xrightarrow{d^V} & \cdots & P_{n-1}^V & \longrightarrow & P_n^V & \longrightarrow & 0 \\
 \text{degree} = & & & \uparrow n & & \uparrow n-1 & & & & & & & \uparrow -n
 \end{array}$$

Since all the boundaries and cycles of P_\bullet and P_\bullet^* are objects of \mathcal{E} , so are $H_i(M_\bullet)$ and $H_i(M_\bullet)^*$. Therefore, M_\bullet and M_\bullet^* are objects of $D_{\mathcal{E}}^b(\mathcal{E})$.

Let $Z_\bullet = \text{cone}(z)$. In [B3, Lemma 3.9], it is further shown that there is a symmetric form $\chi : Z_\bullet \rightarrow Z_\bullet^*$ and that (Z_\bullet, χ) is isometric to $(P_\bullet, s) \perp (Q_\bullet, t)$ in $D^b(\mathcal{E})$. But this tells us that $H_i(Z_\bullet) \cong H_i(P_\bullet) \oplus H_i(Q_\bullet)$ and hence Z_\bullet is an object of $D_{\mathcal{E}}^b(\mathcal{E})$.

Now again following the proof of [B3, Lemma 3.9], it is shown that $T^{-1}z^\# = z$ in $D_{\mathcal{E}}^b(\mathcal{E})$ and that the form χ actually fits in to make M_\bullet a lagrangian for (Z_\bullet, χ) . Hence, this proves the lemma when $n \geq 2$.

In the case $n = 1$, (P_\bullet, s) is given by

$$\begin{array}{ccccccc}
 P_\bullet = & 0 & \longrightarrow & P_1 & \longrightarrow & P_0 & \longrightarrow & P_{-1} & \longrightarrow & 0 \\
 s \downarrow & & & s \downarrow & & s \downarrow & & s \downarrow & & \\
 P_\bullet^* = & 0 & \longrightarrow & P_{-1}^\vee & \longrightarrow & P_0^\vee & \longrightarrow & P_1^\vee & \longrightarrow & 0
 \end{array}$$

Define (Q_\bullet, s) as follows

$$\begin{array}{ccccccc}
 Q_\bullet = \cdots 0 & \longrightarrow & P_1 & \xrightarrow{\begin{pmatrix} s \\ d \end{pmatrix}} & P_{-1}^\vee \oplus P_0 & \longrightarrow & 0 & \longrightarrow & 0 \\
 t \downarrow & & \downarrow & & \downarrow \begin{pmatrix} 0 & d \\ d^\vee & -s \end{pmatrix} & & \downarrow & & \\
 Q_\bullet^* = \cdots 0 & \longrightarrow & 0 & \longrightarrow & P_{-1} \oplus P_0^\vee & \xrightarrow{\begin{pmatrix} s^\vee & d^\vee \end{pmatrix}} & P_1^\vee & \longrightarrow & 0
 \end{array}$$

The degree zero term is in the middle. In [B3, Lemma 3.9], it is established that t is a quasi-isomorphism. It follows that $H_i(Q_\bullet) = 0$ for all $i \neq 0$ and

$$H_0(Q_\bullet) = \frac{P_{-1}^\vee \oplus P_0}{P_1} \in \mathcal{E}.$$

So, Q_\bullet satisfies all the condition in (A.3), because the remaining three conditions are established in [B3, Lemma 3.9]. Now $(P, s) \perp (Q, t)$ has a lagrangian, namely

$$\begin{array}{ccccccc}
 T^{-1}M_\bullet^* = & 0 & \longrightarrow & P_1 & \xrightarrow{-d} & P_0 & \longrightarrow & 0 \\
 z \downarrow & & & \downarrow 0 & & \downarrow sd & & \\
 M_\bullet = & 0 & \longrightarrow & P_0^\vee & \xrightarrow{d^\vee} & P_1^\vee & \longrightarrow & 0 \\
 \uparrow & & & \uparrow & & \uparrow & & \\
 \vdots & & & \vdots & & \vdots & & \\
 \vdots & & & \vdots & & \vdots & & \\
 \text{degree} = & 1 & & 0 & & -1 & &
 \end{array}$$

Again,

$$H_0(M_\bullet) = \ker(d^\vee), \quad H_1(M_\bullet) = \text{coker}(d^\vee) \quad \text{are in } \mathcal{E}.$$

So, M_\bullet and hence M_\bullet^* are objects of $D_{\mathcal{E}}^b(\mathcal{E})$. The rest of the proof is exactly the same as in the case $n \geq 2$. The proof is complete. \square

Finishing the prof of (A.1): We use (A.3) to represent any element x in $W(D_{\mathcal{A}}^b(\mathcal{A}))$ by a chain complex in $D_{\mathcal{A}}^b(\mathcal{A})$ and a strongly symmetric

quasi-isomorphism to its dual. Then, by alternate use of lemma (A.4) and (A.5), we reduce any element in $W(D_{\mathcal{E}}^b(\mathcal{E}))$ to a chain complex in $D_{\mathcal{E}}^b(\mathcal{E})$, concentrated at degree zero. Of course that means the quasi-isomorphism is actually an isomorphism and hence x is the image of an element in $W(\mathcal{A})$ via $W(\mu)$. So $W(\mu)$ is also surjective. So, the proof of theorem (A.1) is complete. \square

References

- [BM] Barge Jean and Morel Fabien, Groupe de Chow des cycles orientés et classe d'Euler des fibrés vectoriels, (French) [The Chow group of oriented cycles and the Euler class of vector bundles], *C. R. Acad. Sci. Paris Sér. I Math.*, **330** (2000) no. 4, 287–290.
- [B1] Balmer Paul, Derived Witt groups of a scheme, *J. Pure Appl. Algebra*, **141** (1999) no. 2, 101–129.
- [B2] Balmer Paul, Triangular Witt groups Part I: The 12-term localization exact sequence, *K-Theory*, **19** (2000) 311–363.
- [B3] Balmer Paul, Triangular Witt groups. II. From usual to derived, *Math. Z.*, **236** (2001) no. 2, 351–382.
- [B4] Balmer Paul, Witt cohomology, Mayer-Vietoris, homotopy invariance and the Gersten conjecture, *K-Theory*, **23** (2001) no. 1, 15–30.
- [BGPW] Balmer Paul, Gille Stefan, Panin Ivan and Walter Charles, The Gersten conjecture for Witt groups in the equicharacteristic case, *Doc. Math.*, **7** (2002) 203–217.
- [BW] Balmer Paul and Walter Charles, A Gersten-Witt spectral sequence for regular schemes, *Ann. Sci. École Norm. Sup. (4)*, **35** (2002) no. 1, 127–152.
- [CS] Colliot-Thélène Jean-Louis and J.-J. Sansuc, Fibrés quadratiques et composantes connexes réelles, (French) *Math. Ann.*, **244** (1979) no. 2, 105–134.
- [F1] Fasel Jean, Groupes de Chow-Witt, *Mém. Soc. Math. Fr. (N.S.)*, **113** (2008).
- [F2] Fasel Jean, The Chow-Witt ring, *Documenta Mathematica*, **12** (2007) 275–312.
- [FS] J. Fasel and V. Srinivas, Chow-Witt groups and Grothendieck-Witt groups of regular schemes, *Advances in Mathematics*, **221** (2009) 302–329.
- [G1] S. Gille, On Witt groups with support, *Math. Ann.*, **322** (2002) no. 1, 103–137.
- [G2] S. Gille, A transfer morphism for Witt groups, *J. Reine Angew. Math.*, **564** (2003) 215–233.
- [K] Keller Bernhard, On the cyclic homology of exact categories, *J. Pure Appl. Algebra*, **136** (1999) no. 1, 1–56.
- [M1] Satya Mandal, Sublagrangian constructions, *J. Ramanujan Math. Soc.*, **30** (2015) no. 3, 309–330.
- [M2] Satya Mandal, Derived Witt group formalism, *J. Pure Appl. Algebra*, **219** (2015) no. 8, 3518–3534.
- [MS] Satya Mandal and Sarang Sane, Stability of locally CMFPD homologies under duality, *Journal of Algebra*, **440** (2015) 49–71. Preprint.
- [Mo] F. Morel, \mathbb{A}^1 -algebraic topology over a field, LNM 2052, Springer, Heidelberg (2012).
- [OP] Ojanguren Manuel and Panin Ivan, A purity theorem for the Witt group, *Ann. Sci. École Norm. Sup. (4)*, **32** (1999) no. 1, 71–86.
- [OPSS] M. Ojanguren, R. Parimala, R. Sridharan and V. Suresh, Witt groups of the punctured spectrum of a 3-dimensional regular local ring and a purity theorem, *J. London Math. Soc. (2)*, **59** (1999) no. 2, 521–540.

- [Pa1] Pardon William, A “Gersten conjecture” for Witt groups, Algebraic K-theory, Part II (Oberwolfach, 1980), 300–315, Lecture Notes in Math., Springer, Berlin, **967** (1982).
- [Pa2] Pardon William, A relation between Witt groups and zero-cycles in a regular ring, Algebraic K-theory, number theory, geometry and analysis (Bielefeld, 1982), 261–328, Lecture Notes in Math., Springer, Berlin, **1046** (1984).
- [Pa3] William Pardon, The filtered Gersten-Witt resolution for regular schemes, <http://www.math.uiuc.edu/K-theory/0419/>.
- [QSS] H.-G. Quebbemann, W. Scharlau and M. Schulte, Quadratic and Hermitian forms in additive and abelian categories, *J. Algebra*, **59** (1979) no. 2, 264–289.
- [W] A. Weibel Charles, An introduction to homological algebra, Cambridge Studies in Advanced Mathematics, 38. Cambridge University Press, Cambridge (1994) pp. xiv+450.